

Local controllability to trajectories of the magnetohydrodynamic equations

Mehdi Badra

Abstract. In this paper we prove the local controllability to trajectories of the three dimensional magnetohydrodynamic equations by means of two internal controls, one in the velocity equations and the other in the magnetic field equations and both localized in an arbitrary small subset with not empty interior of the domain. This paper improves the previous results [6, 7, 31, 32] where the second control is not localized and it allows to deduce the local controllability to trajectories with boundary controls. The proof relies on the Carleman inequality for the Stokes system of [36] to deal with the velocity equations and on a new Carleman inequality for a Dynamo-type equation to deal with the magnetic field equations.

Mathematics Subject Classification (2010). Primary 76W05, 93B05; Secondary 76D55, 35Q35.

Keywords. magnetohydrodynamic equations, null controllability, Carleman inequality.

1. Introduction

In the present paper we are interested in proving the local controllability to trajectories of the following magnetohydrodynamic (MHD) equations:

$$\left\{ \begin{array}{ll} v_t - \Delta v + (v \cdot \nabla)v - (\operatorname{curl} B) \times B + \nabla p = \bar{f} + \mathbf{1}_O u_1 & \text{in } Q, \\ B_t + \operatorname{curl}(\operatorname{curl} B) - \operatorname{curl}(v \times B) = \mathbf{1}_O u_2 & \text{in } Q, \\ \operatorname{div} v = \operatorname{div} B = 0 & \text{in } Q, \\ v = \bar{v}, \quad B \cdot n = \bar{h} & \text{on } \Sigma, \\ (\operatorname{curl} B - v \times B) \times n = \bar{e} & \text{on } \Sigma, \\ v(0) = v_0, \quad B(0) = B_0 & \text{in } \Omega. \end{array} \right. \quad (1.1)$$

The system (1.1) describes the motion of an incompressible and electrically conducting fluid in a bounded domain Ω of \mathbb{R}^3 during the time interval $(0, T)$ where T is a fixed positive time horizon. Above and in the following \times denotes

the vectorial product, $n(x) \in \mathbb{R}^3$ denotes the unit exterior normal vector field defined on the boundary $\partial\Omega$ which is supposed to be regular. Moreover, we use the notations $Q \stackrel{\text{def}}{=} (0, T) \times \Omega$ and $\Sigma \stackrel{\text{def}}{=} (0, T) \times \partial\Omega$ for cylindrical domains.

The fluid is characterized by its velocity $v(t, x) \in \mathbb{R}^3$, by a pressure function $p(t, x) \in \mathbb{R}$ and by the magnetic field $B(t, x) \in \mathbb{R}^3$. The system (1.1) is a coupling between the incompressible Navier-Stokes equations satisfied by the velocity and the induction equations satisfied by the magnetic field. These last are obtained from the Maxwell's equations and the coupling is due to Lorentz forces through the presence of terms $(\text{curl } B) \times B$ and $v \times B$, we refer to [13] for a detailed explanation of the derivation of (1.1). Note that here, the usual non-dimensional constants that characterize the flow (Hartmann number, interaction parameter and magnetic Reynolds number) are supposed to be equal to one for simplicity.

In (1.1), $\bar{f}(t, x) \in \mathbb{R}^3$ is a given body force and $\bar{v}(t, x) \in \mathbb{R}^3$, $\bar{h}(t, x) \in \mathbb{R}$, $\bar{e}(t, x) \in \mathbb{R}^3$ are the boundary values of a given target solution (\bar{v}, \bar{B}) which satisfies:

$$\left\{ \begin{array}{ll} \bar{v}_t - \Delta \bar{v} + (\bar{v} \cdot \nabla) \bar{v} - (\text{curl } \bar{B}) \times \bar{B} + \nabla \bar{p} = \bar{f} & \text{in } Q, \\ \bar{B}_t + \text{curl}(\text{curl } \bar{B}) - \text{curl}(\bar{v} \times \bar{B}) = 0 & \text{in } Q, \\ \text{div } \bar{v} = \text{div } \bar{B} = 0 & \text{in } Q, \\ \bar{B} \cdot n = \bar{h}, \quad (\text{curl } \bar{B} - \bar{v} \times \bar{B}) \times n = \bar{e} & \text{on } \Sigma. \end{array} \right. \quad (1.2)$$

To be compatible with induction and divergence equalities in (1.2) these boundary values must satisfy:

$$\begin{aligned} \int_{\partial\Omega} \bar{v} \cdot n d\Gamma &= \int_{\partial\Omega} \bar{h} d\Gamma = 0 \quad \text{on } (0, T), \\ \bar{h}_t &= \text{div}_\tau \bar{e} \quad \text{on } \Sigma, \end{aligned} \quad (1.3)$$

where div_τ is the surfacic divergence operator. The second line in (1.3) is obtained by taking the normal trace of the magnetic field equation. Note that the particular situation where $\bar{h} = 0$ and $\bar{e} = 0$ corresponds to a perfectly conductive boundary.

In (1.1), u_1, u_2 are control functions in $\mathbf{L}^2((0, T) \times \mathcal{O})$ which are used to reach the target state (\bar{v}, \bar{B}) at time T . Here \mathcal{O} is a nonempty and simply connected open subset of Ω , $\mathbf{1}_{\mathcal{O}} : \mathbf{L}^2(\mathcal{O}) \rightarrow \mathbf{L}^2(\Omega)$ denotes the extension operator defined by $\mathbf{1}_{\mathcal{O}}(z)(x) = z(x)$ if $x \in \mathcal{O}$ and $\mathbf{1}_{\mathcal{O}}(z)(x) = 0$ else. Note that, above and in the following, we write in bold the spaces of vector-valued functions: $\mathbf{L}^2(\Omega) = (L^2(\Omega))^3$, $\mathbf{L}^2(\mathcal{O}) = (L^2(\mathcal{O}))^3$ etc. We underline that to guarantee the well-posedness of the controlled induction equations without an additional gradient of pressure we need to search a control u_2 satisfying

$$\begin{aligned} \text{div}(\mathbf{1}_{\mathcal{O}} u_2) &= 0 \quad \text{in } Q, \\ (\mathbf{1}_{\mathcal{O}} u_2) \cdot n &= 0 \quad \text{on } \Sigma. \end{aligned} \quad (1.4)$$

For that, we choose u_2 of the form

$$u_2 = P_{\mathcal{O}} \tilde{u}_2 \quad (1.5)$$

where $\tilde{u}_2 \in \mathbf{L}^2((0, T) \times \mathcal{O})$ and $P_{\mathcal{O}}$ is the classical Helmholtz projector related to \mathcal{O} (i.e the orthogonal projection operator from $\mathbf{L}^2(\mathcal{O})$ onto the completion of $\{v \in \mathbf{C}_c^\infty(\mathcal{O}) \mid \operatorname{div} v = 0 \text{ in } \mathcal{O}\}$ for the norm of $\mathbf{L}^2(\mathcal{O})$).

System (1.1) is said to be controllable to trajectories in time T , if for a given target solution (\bar{v}, \bar{B}) satisfying (1.2) and a given pair of initial data (v_0, B_0) there exist controls u_1, u_2 such that the solution (v, B) of (1.1) satisfies the terminal conditions:

$$v(T) = \bar{v}(T) \quad \text{and} \quad B(T) = \bar{B}(T) \quad \text{in } \Omega. \quad (1.6)$$

Such a property is said to be local if it holds for initial data v_0, B_0 close enough to $\bar{v}(0), \bar{B}(0)$. It is this last property that we intend to prove in this paper.

From a practical point of view, the controllability to trajectories is an important notion because it implies the (open loop) stabilizability which is the first property to be checked in order to construct a stabilizing feedback control. For the stabilization of a parabolic system around a stationary state, a criterion weaker than exact controllability is known (see [5]) but if the target state is not stationary, prove the exact controllability seems to be the most reasonable alternative. For instance in [8], the design of a feedback control stabilizing the Navier-Stokes equations around an instationary state is deduced from the controllability properties of the Navier-Stokes equations.

Before to state our result, let us clarify our hypotheses. We assume that

$$\begin{aligned} \partial\Omega \text{ is a two dimensional manifold of class } C^{2,1} \text{ and is} \\ \text{composed of a finite number of connected components,} \end{aligned} \quad (1.7)$$

and that

$$\begin{aligned} \bar{v} &\in L^2(0, T; \mathbf{H}^2(\Omega)) \cap H^1(0, T; \mathbf{L}^2(\Omega)), \\ \bar{B} &\in L^2(0, T; \mathbf{H}^2(\Omega)) \cap H^1(0, T; \mathbf{L}^2(\Omega)). \end{aligned} \quad (1.8)$$

We also assume the following compatibility conditions for the initial data:

$$\begin{aligned} \operatorname{div} v_0 = \operatorname{div} B_0 = 0 \quad \text{in } \Omega \\ (v_0 - \bar{v}(0)) \cdot n = (B_0 - \bar{B}(0)) \cdot n = 0 \quad \text{on } \partial\Omega. \end{aligned} \quad (1.9)$$

Next, we denote by $g_i, i = 1, \dots, N$ a basis of the finite dimensional space:

$$X_N \stackrel{\text{def}}{=} \{y \in \mathbf{L}^2(\Omega) \mid \operatorname{div} y = 0 \text{ in } \Omega, \operatorname{curl} y = 0 \text{ in } \Omega, y \cdot n = 0 \text{ on } \partial\Omega\}. \quad (1.10)$$

The fact that the above space is finite dimensional is well-known, see for instance [12, Chap. IX] or [19] for a detailed characterization of X_N . We recall that if Ω is simply-connected we have $N = 0$ and X_N reduces to $\{0\}$ and if Ω is multiply-connected then $N \geq 1$ is the number of cuts required to make Ω simply-connected. Moreover, since \mathcal{O} is simply connected we have $g_i = \nabla p_i$ in \mathcal{O} for some smooth function p_i and with (1.5) it yields:

$$\int_{\Omega} \mathbf{1}_{\mathcal{O}} u_2 \cdot g_i dx = \int_{\mathcal{O}} P_{\mathcal{O}} \tilde{u}_2 \cdot \nabla p_i dx = 0. \quad (1.11)$$

As a consequence, if we multiply the second equality in (1.1) and in (1.2) by g_i and integrate over Ω , with an integration by parts we obtain

$$\frac{d}{dt} \int_{\Omega} B(t) \cdot g_i dx = - \int_{\partial\Omega} \bar{e}(t) \cdot g_i d\Gamma = \frac{d}{dt} \int_{\Omega} \bar{B}(t) \cdot g_i dx \quad (1.12)$$

which means that $\int_{\Omega} (B(t) - \bar{B}(t)) \cdot g_i dx$ must be constant. Then for the following we make the additional assumption

$$\forall i = 1, \dots, N \quad \int_{\Omega} (B_0 - \bar{B}(0)) \cdot g_i dx = 0, \quad (1.13)$$

which implies the following condition for the magnetic field:

$$\forall i = 1, \dots, N \quad \int_{\Omega} B(t) \cdot g_i dx = \int_{\Omega} \bar{B}(t) \cdot g_i dx \quad \text{for a.e. } t \in (0, T). \quad (1.14)$$

Note that if we denote by C_i , $i = 1, \dots, N$ a family of smooth cuts required to make Ω simply-connected then it is well-known (see [19, Lemme 1.4]) that (1.13) and (1.14) are respectively equivalent to $\int_{C_i} (B_0 - \bar{B}(0)) \cdot nd\Gamma = 0$ and

$$\int_{C_i} B(t) \cdot nd\Gamma = \int_{C_i} \bar{B}(t) \cdot nd\Gamma \quad \forall t \in (0, T), \quad \forall i = 1, \dots, N.$$

We are now in position to state our main theorem. In what follows, $\mathbf{H}_{00}^{1/2}(\Omega)$ stands for the subspace of $\mathbf{H}_0^{1/2}(\Omega)$ composed with y such that $\int_{\Omega} \text{dist}(x, \partial\Omega)^{-1} |y|^2 dx < +\infty$.

Theorem 1.1. *Assume that Ω is a bounded domain of \mathbb{R}^3 and that (1.7), (1.8) are satisfied. There exists $c_0 > 0$ and $\varepsilon > 0$ such that for all $(v_0, B_0) \in \mathbf{H}_{00}^{1/2}(\Omega) \times \mathbf{H}^{1/2}(\Omega)$ satisfying (1.9), (1.13) and*

$$\|(v_0 - \bar{v}(0), B_0 - \bar{B}(0))\|_{\mathbf{H}_{00}^{1/2}(\Omega) \times \mathbf{H}^{1/2}(\Omega)} \leq \varepsilon$$

there exists $(u_1, u_2) \in \mathbf{L}^2((0, T) \times \mathcal{O}) \times \mathbf{L}^2((0, T) \times \mathcal{O})$ satisfying (1.4) such that system (1.1), (1.14) admits a solution

$$(v, B) \in L^2(0, T; \mathbf{H}^{3/2}(\Omega) \times \mathbf{H}^{3/2}(\Omega)) \cap C(0, T; \mathbf{H}^{1/2}(\Omega) \times \mathbf{H}^{1/2}(\Omega))$$

which satisfies the terminal condition (1.6) as well as the estimate:

$$\begin{aligned} & \| (v - \bar{v}, B - \bar{B}) \|_{L^2(0, T; \mathbf{H}^{3/2}(\Omega) \times \mathbf{H}^{3/2}(\Omega))} + \| (v - \bar{v}, B - \bar{B}) \|_{C(0, T; \mathbf{H}_{00}^{1/2}(\Omega) \times \mathbf{H}^{1/2}(\Omega))} \\ & \quad + \| (u_1, u_2) \|_{\mathbf{L}^2((0, T) \times \mathcal{O}) \times \mathbf{L}^2((0, T) \times \mathcal{O})} \\ & \leq c_0 \| (v_0 - \bar{v}(0), B_0 - \bar{B}(0)) \|_{\mathbf{H}_{00}^{1/2}(\Omega) \times \mathbf{H}^{1/2}(\Omega)}. \end{aligned}$$

Moreover, if $(\tilde{v}, \tilde{B}) \in L^2(0, T; \mathbf{H}^{3/2}(\Omega) \times \mathbf{H}^{3/2}(\Omega)) \cap C(0, T; \mathbf{H}^{1/2}(\Omega) \times \mathbf{H}^{1/2}(\Omega))$ is another solution of (1.1) corresponding to the initial data v_0, B_0 and to the controls u_1, u_2 then (\tilde{v}, \tilde{B}) and (v, B) must coincide i.e $(\tilde{v}, \tilde{B}) \equiv (v, B)$.

Remark 1.2. 1. The main novelty of the above controllability result is that the two controls are localized in a small subset of the domain. This improves the results of [6, 7, 31, 32] where only the control in the velocity equation is localized.

2. The regularity assumptions (1.8) are weaker than those made in previous work on the subject, see in particular [31, 32]. Moreover, the initial regularity assumption $(v_0 - \bar{v}(0), B_0 - \bar{B}(0)) \in \mathbf{H}_{00}^{1/2}(\Omega) \times \mathbf{H}^{1/2}(\Omega)$ is “optimal” in the sense that it is the weaker as possible to obtain a fixed-point solution as well as its uniqueness.
3. Although it follows from standard arguments, the second part of the theorem about the uniqueness of the controlled solution must be underlined because it guarantees that the action of the control is effective: there does not exist another “uncontrolled” trajectory (v, B) corresponding to internal body forces u_1, u_2 and which does not satisfy (1.6). Note that such a uniqueness property is not clear when dealing with density dependent fluid, see the forthcoming work [4].
4. If (1.13) is not satisfied then the terminal condition $B(T) = \bar{B}(T)$ cannot be obtained because (1.12) would imply that the equalities in (1.14) are violated at $t = T$. However, it is possible to get rid of (1.13) by adding a global control term in the induction equations. More precisely, an analogue version of Theorem 1.1 can be obtained without the assumption (1.13) and for a control with a global gradient term of the form

$$\mathbf{1}_{\mathcal{O}} u_2 + \nabla \chi, \quad \chi \in L^2(0, T; H^1(\Omega)),$$

or for a control with a finite dimensional global term of the form

$$\mathbf{1}_{\mathcal{O}} P_{\mathcal{O}} u_2 + \sum_{i=1}^N \alpha_i g_i, \quad \alpha_i \in L^2(0, T), \quad i = 1, \dots, N.$$

The adaptation of the proof is left to the reader.

The two dimensional counterpart of Theorem 1.1 is also true. If Ω is supposed to be only bounded in directions x_1, x_2 and invariant in direction x_3 , if all functions only depend on x_1, x_2 and if each vector field has its third component equal to zero (and then is identified to its two first components i.e $v(x) = {}^t(v_1(x_1, x_2), v_2(x_1, x_2), 0) \equiv {}^t(v_1(x_1, x_2), v_2(x_1, x_2))$, etc), then we can identify Ω to its bounded two dimensional section and (1.1) reduces to:

$$\left\{ \begin{array}{ll} v_t - \Delta v + (v \cdot \nabla)v + (\operatorname{curl} B)B^\perp + \nabla p = \bar{f} + \mathbf{1}_{\mathcal{O}} u_1 & \text{in } Q, \\ B_t + \operatorname{curl}(\operatorname{curl} B) - \operatorname{curl}(v \cdot B^\perp) = \mathbf{1}_{\mathcal{O}} u_2 & \text{in } Q, \\ \operatorname{div} v = \operatorname{div} B = 0 & \text{in } Q, \\ v = \bar{v}, \quad B \cdot n = \bar{h} & \text{on } \Sigma, \\ (\operatorname{curl} B - v \cdot B^\perp) = \bar{e} & \text{on } \Sigma, \\ v(0) = v_0, \quad B(0) = B_0 & \text{in } \Omega, \end{array} \right. \quad (1.15)$$

where we have used the notations ${}^t(a_1, a_2)^\perp \stackrel{\text{def}}{=} {}^t(a_2, -a_1)$, $\operatorname{curl} a \stackrel{\text{def}}{=} \partial_{x_1} a_2 - \partial_{x_2} a_1$ for a vector field $a = {}^t(a_1, a_2)$ and $\operatorname{curl} b \stackrel{\text{def}}{=} {}^t(\partial_{x_2} b, -\partial_{x_1} b)$ for a scalar function b . Then the following two dimensional version of Theorem 1.1 holds.

Theorem 1.3. *Assume that Ω is a bounded domain of \mathbb{R}^2 and that (1.7), (1.8) are satisfied. There exists $c_0 > 0$ and $\varepsilon > 0$ such that for all $(v_0, B_0) \in \mathbf{L}^2(\Omega) \times \mathbf{L}^2(\Omega)$ satisfying (1.9), (1.13) and*

$$\|(v_0 - \bar{v}(0), B_0 - \bar{B}(0))\|_{\mathbf{L}^2(\Omega) \times \mathbf{L}^2(\Omega)} \leq \varepsilon$$

there exists $(u_1, u_2) \in \mathbf{L}^2((0, T) \times \mathcal{O}) \times \mathbf{L}^2((0, T) \times \mathcal{O})$ satisfying (1.4) such that system (1.15), (1.14) admits a solution $(v, B) \in L^2(0, T; \mathbf{H}^1(\Omega) \times \mathbf{H}^1(\Omega)) \cap C(0, T; \mathbf{L}^2(\Omega) \times \mathbf{L}^2(\Omega))$ which satisfies the terminal condition (1.6) as well as estimate

$$\begin{aligned} & \|(v - \bar{v}, B - \bar{B})\|_{L^2(0, T; \mathbf{H}^1(\Omega) \times \mathbf{H}^1(\Omega))} + \|(v - \bar{v}, B - \bar{B})\|_{C(0, T; \mathbf{L}^2(\Omega) \times \mathbf{L}^2(\Omega))} \\ & \quad + \|(u_1, u_2)\|_{\mathbf{L}^2((0, T) \times \mathcal{O}) \times \mathbf{L}^2((0, T) \times \mathcal{O})} \\ & \leq c_0 \|(v_0 - \bar{v}(0), B_0 - \bar{B}(0))\|_{\mathbf{L}^2(\Omega) \times \mathbf{L}^2(\Omega)}. \end{aligned}$$

Moreover, if $(\tilde{v}, \tilde{B}) \in L^2(0, T; \mathbf{H}^1(\Omega) \times \mathbf{H}^1(\Omega)) \cap C(0, T; \mathbf{L}^2(\Omega) \times \mathbf{L}^2(\Omega))$ is another solution of (1.15) corresponding to the initial data v_0, B_0 and to the controls u_1, u_2 then (\tilde{v}, \tilde{B}) and (v, B) must coincide i.e $(\tilde{v}, \tilde{B}) \equiv (v, B)$.

The proof of Theorem 1.3 is simpler than the one of Theorem 1.1, see Remark 4.4 below. Then in the following we omit it and we only consider the three dimensional case.

The local controllability of magnetohydrodynamic equations has been studied in [6, 7, 31, 32]. However, these quoted works provide controllability results with a non local control of the form $P\mathbf{1}_{\mathcal{O}}u_2$ in the induction equations where u_2 does not necessarily satisfies (1.4) and where P is the Helmholtz projector related to Ω . This means that the control is of the form $\mathbf{1}_{\mathcal{O}}u_2 + \nabla\chi$ in Q for some pressure function χ not necessarily supported in \mathcal{O} . As a consequence, these results cannot be used to deduce the controllability with a boundary control from an extension of the domain procedure. Such a drawback has already been pointed out in [39, 5] when considering the feedback stabilizability of magnetohydrodynamic equations around a stationary solution. To deduce from above theorems the local controllability to trajectories with boundary control localized in an arbitrary small subset $\Gamma_c \subset \partial\Omega$, it suffices to extend Ω through Γ_c to some domain $\tilde{\Omega}$ and to apply Theorem 1.1 (or Theorem 1.3) to the MHD system extended to $\tilde{\Omega}$ for an internal control localized in $\mathcal{O} \subset \tilde{\Omega} \setminus \Omega$. The boundary control will then be obtained by taking the restriction to $(0, T) \times \Gamma_c$ of the controlled solution defined in $(0, T) \times \tilde{\Omega}$. We then obtain control functions (b_1, b_2, b_3) supported in Γ_c and satisfying

$$\begin{aligned} \int_{\partial\Omega} b_1 \cdot n d\Gamma &= \int_{\partial\Omega} b_2 d\Gamma = 0 \quad \text{on } (0, T), \\ (b_2)_t &= \text{div}_\tau b_3 \quad \text{on } \Sigma, \end{aligned} \tag{1.16}$$

such that the conclusion of Theorem 1.1 (or of Theorem 1.3) remains true for system (1.1) with $u_1 = u_2 = 0$ and with the boundary data replaced by

$$\begin{aligned} v &= \bar{v} + b_1, & B \cdot n &= \bar{h} + b_2 & \text{on } \Sigma, \\ (\text{curl } B - v \times B) \times n &= \bar{e} + b_3 & \text{on } \Sigma. \end{aligned}$$

Here we follow a classical approach for proving the controllability of a parabolic system which consists in proving a global Carleman inequality for an adjoint system. It is a type of sophisticated weighted inequality which permits to bound the values of the solution in the whole domain in term of its values in a subdomain. Global Carleman inequalities were first introduced in [22, 23, 24] and then were extensively used to prove the exact controllability of Stokes-type system. About the exact controllability of Stokes and Navier-Stokes equations we refer to [22, 20, 10, 21, 34, 17, 18, 29, 11] about some other related coupled systems we refer to [28, 37, 26, 30, 16, 41]. We must underline that our result deeply relies on a Carleman inequality for the Stokes system which has been recently proved in [36] (see also [35, 3]).

To prove Theorem 1.1 we first remark that the local controllability to trajectories of (1.1), (1.14) is equivalent to the local null controllability of the system satisfied by $(w, \theta) = (v - \bar{v}, B - \bar{B})$. By taking (1.5) into account and renaming \tilde{u}_2 by u_2 for simplicity, the problem is then reduced to find (u_1, u_2) such that the solution (w, B) of

$$\left\{ \begin{array}{ll} w_t - \Delta w + (w \cdot \nabla) \bar{v} + (\bar{v} \cdot \nabla) w - (\bar{B} \cdot \nabla) \theta \\ \quad - (\theta \cdot \nabla) \bar{B} - (\theta \cdot \nabla) \theta + (w \cdot \nabla) w + \nabla q = \mathbf{1}_{\mathcal{O}} u_1 & \text{in } Q, \\ \theta_t + \text{curl}(\text{curl} \theta) - \text{curl}(w \times \bar{B} + \bar{v} \times \theta) \\ \quad - \text{curl}(w \times \theta) = \mathbf{1}_{\mathcal{O}} P_{\mathcal{O}} u_2 & \text{in } Q, \\ \text{div } w = \text{div } \theta = 0 & \text{in } Q, \\ \forall i = 1, \dots, N \quad \int_{\Omega} w \cdot g_i dx = 0 & \text{in } (0, T), \\ w = 0, \quad \theta \cdot n = 0 & \text{on } \Sigma, \\ (\text{curl} \theta - \bar{v} \times \theta) \times n = 0 & \text{on } \Sigma, \\ w(0) = v_0 - \bar{v}(0), \quad \theta(0) = B_0 - \bar{B}(0) & \text{in } \Omega, \end{array} \right. \quad (1.17)$$

satisfies

$$w(T) = 0 \quad \text{and} \quad \theta(T) = 0 \quad \text{in } \Omega. \quad (1.18)$$

Note that to obtain that $(w, \theta) = (v - \bar{v}, B - \bar{B})$ satisfies (1.17) we have used the formula of vectorial analysis $(\text{curl } a) \times a = (a \cdot \nabla) a - \nabla(|a|^2/2)$. In order to construct a solution to (1.17)-(1.18) with a fixed point procedure, a first step consists in replacing the nonlinear terms $(\theta \cdot \nabla) \theta - (w \cdot \nabla) w$ and $\text{curl}(w \times \theta)$ by adequate nonhomogeneous terms g_1 and g_2 vanishing at time T and to

construct a solution of the corresponding linear system:

$$\left\{ \begin{array}{ll} w_t - \Delta w + (w \cdot \nabla)\bar{v} + (\bar{v} \cdot \nabla)w - (\bar{B} \cdot \nabla)\theta & \\ \quad \quad \quad -(\theta \cdot \nabla)\bar{B} + \nabla q = g_1 + \mathbf{1}_O u_1 & \text{in } Q, \\ \theta_t + \text{curl}(\text{curl } \theta) - \text{curl}(w \times \bar{B} + \bar{v} \times \theta) = g_2 + \mathbf{1}_O P_O u_2 & \text{in } Q, \\ \quad \quad \quad \text{div } w = \text{div } \theta = 0 & \text{in } Q, \\ \forall i = 1, \dots, N \quad \int_{\Omega} w \cdot g_i dx = 0 & \text{in } (0, T), \quad (1.19) \\ \quad \quad \quad w = 0, \quad \theta \cdot n = 0 & \text{on } \Sigma, \\ \quad \quad \quad (\text{curl } \theta - \bar{v} \times \theta) \times n = 0 & \text{on } \Sigma, \\ \quad \quad \quad w(0) = w_0, \quad \theta(0) = \theta_0 & \text{in } \Omega, \\ \quad \quad \quad w(T) = 0, \quad \theta(T) = 0 & \text{in } \Omega. \end{array} \right.$$

For that, we use a classical duality argument which relies on a Carleman inequality for the adjoint system:

$$\left\{ \begin{array}{ll} -y_t - \Delta y - (D^s y)\bar{v} + (D^a \rho)\bar{B} + \nabla \pi = f_1 & \text{in } Q, \\ -\rho_t - \Delta \rho + (D^s y)\bar{B} - (D^a \rho)\bar{v} + \nabla \kappa + \sum_{i=1}^N \mu_i g_i = f_2 & \text{in } Q, \\ \quad \quad \quad \text{div } y = \text{div } \rho = 0 & \text{in } Q, \quad (1.20) \\ \forall i = 1, \dots, N \quad \int_{\Omega} \rho \cdot g_i dx = 0 & \text{in } (0, T), \\ \quad \quad \quad y = 0, \quad \rho \cdot n = 0, \quad (\text{curl } \rho) \times n = 0 & \text{on } \Sigma, \end{array} \right.$$

where we have used the notations $D^s y \stackrel{\text{def}}{=} \nabla y + {}^t \nabla y$ and $D^a y \stackrel{\text{def}}{=} \nabla y - {}^t \nabla y$. In (1.20), the pressure functions π and κ and the real values μ_i , $i = 1, \dots, N$ plays the role of the Lagrange multipliers related to the free divergence constraints and to conditions $\int_{\Omega} \rho \cdot g_i dx = 0$, $i = 1, \dots, N$.

To obtain a Carleman inequality for (1.20) we apply to the first equality the Carleman inequality for the Stokes system of [36] and we apply to the second equality a new Carleman inequality for a Dynamo-type equation that we prove in Section 4. We underline that unlike in [6, 7, 31, 32] we do not treat directly the Dynamo-type equation but we prove a Carleman inequality for the curl of such equation, which has the advantage of getting rid of the terms $\nabla \kappa$ and $\mu_i g_i$. This last inequality is obtained from a Carleman inequality for a vectorial heat equation with an homogeneous Robin boundary condition on the normal component and with an homogeneous Dirichlet boundary condition on the tangential component. For that, we use the idea of [14] to get rid of boundary terms corresponding to the second Carleman parameter with odd exponents.

The rest of the paper is organized as follows. The Section 2 is dedicated to notations. The Section 3 is devoted to the statements of theorems related to each steps of the proof of Theorem 1.1, their proof are postponed to sections 4 and 5. The Section 4 concerns the proof of the Carleman inequality

for the adjoint MHD system (1.20). In particular, we provide a Carleman inequality for a vectorial heat equation with an homogeneous Robin boundary condition on the normal component and with an homogeneous Dirichlet boundary condition on the tangential component. Section 5 is dedicated to more classical proofs: we deduce the existence of a solution of (1.19) from a duality argument and we deduce the local null controllability of the nonlinear MHD systems (1.17) from a fixed-point procedure. Finally, we postpone in an Appendix some regularity results for the Stokes system with a nonhomogeneous non standard boundary condition and for linear non autonomous magnetohydrodynamic equations.

2. Notations and definitions

Let us recall that Ω is a bounded open subset of \mathbb{R}^3 with a regular boundary $\partial\Omega$ satisfying (1.7). We denote by $n = {}^t(n_1, n_2, n_3)$ the outward unit normal to $\partial\Omega$ (t denotes the transpose). For a scalar function w or a vector field $y = {}^t(y_1, y_2, y_3)$ we define $\nabla w \stackrel{\text{def}}{=} {}^t(\partial_{x_1} w, \partial_{x_2} w, \partial_{x_3} w)$, $\nabla y \stackrel{\text{def}}{=} (\partial_{x_j} y_i)_{1 \leq i, j \leq 3}$, ${}^t\nabla y \stackrel{\text{def}}{=} (\partial_{x_i} y_j)_{1 \leq i, j \leq 3}$ and we use the notations $D^s y \stackrel{\text{def}}{=} \nabla y + {}^t\nabla y$ and $D^a y \stackrel{\text{def}}{=} \nabla y - {}^t\nabla y$. Moreover we define the normal derivatives $\frac{\partial w}{\partial n} \stackrel{\text{def}}{=} (\nabla w) \cdot n$ and $\frac{\partial y}{\partial n} \stackrel{\text{def}}{=} (\nabla y)n$ on $\partial\Omega$, we use the notations $y_n \stackrel{\text{def}}{=} y \cdot n$ and $y_\tau \stackrel{\text{def}}{=} y - y_n n$ for the normal and the tangential components of y on $\partial\Omega$ and we recall that $y_\tau = 0$ if and only if $y \times n = 0$ where \times denotes the vectorial product. The divergence of y is defined by $\text{div } y \stackrel{\text{def}}{=} \sum_{j=1}^3 \partial_{x_j} y_j$ and the curl of y is defined by

$$\text{curl } y \stackrel{\text{def}}{=} \begin{pmatrix} \partial_{x_2} y_3 - \partial_{x_3} y_2 \\ \partial_{x_3} y_1 - \partial_{x_1} y_3 \\ \partial_{x_1} y_2 - \partial_{x_2} y_1 \end{pmatrix}.$$

We denote by $L^2(\Omega)$, $L^2(\partial\Omega)$, $H^r(\Omega)$, $H^r(\partial\Omega)$, $H_0^r(\Omega)$ and $H^{-r}(\Omega) = [H_0^r(\Omega)]'$ for $r \geq 0$, the usual Lebesgue and Sobolev spaces of scalar functions in Ω or in $\partial\Omega$, we define $H_{00}^{1/2}(\Omega)$ as the subspace of $H_0^{1/2}(\Omega)$ composed with functions y satisfying $\int_\Omega \text{dist}(x, \partial\Omega)^{-1} |y|^2 dx < +\infty$ (see [40, Thm. 11.7]), and we write in bold the spaces of vector-valued functions: $\mathbf{L}^2(\Omega) = (L^2(\Omega))^3$, $\mathbf{L}^2(\partial\Omega) = (L^2(\partial\Omega))^3$, etc. Above and in the following $[X]'$ stands for the dual space of X . Moreover, we define $\mathbf{H}_n^2(\Omega)$ as the space of functions $y \in \mathbf{H}^2(\Omega)$ such that $y \cdot n = 0$ on $\partial\Omega$ and for $r \in [0, 2]$ we define $\mathbf{H}_n^r(\Omega)$ as the interpolation space $[\mathbf{H}_n^2(\Omega), \mathbf{L}^2(\Omega)]_{1-r/2}$. Standard arguments guarantee that $\mathbf{H}_n^r(\Omega) = \mathbf{H}^r(\Omega)$ if $r \in [0, 1/2)$, that $\mathbf{H}_n^r(\Omega) = \{y \in \mathbf{H}^r(\Omega) \mid y \cdot n = 0 \text{ on } \partial\Omega\}$ if $r \in (1/2, 2]$ and that $\mathbf{H}_n^{1/2}(\Omega) = \{y \in \mathbf{H}^{1/2}(\Omega) \mid y \cdot n \in H_{00}^{1/2}(\Omega)\}$.

We also introduce the spaces of free divergence functions for $r \in [0, 2]$:

$$\mathbf{V}_n^r(\Omega) \stackrel{\text{def}}{=} \left\{ y \in \mathbf{H}^r(\Omega) \mid \text{div } y = 0 \text{ in } \Omega, \quad y \cdot n = 0 \text{ on } \partial\Omega \right\},$$

$$\mathbf{V}_0^r(\Omega) \stackrel{\text{def}}{=} [\mathbf{V}_n^r(\Omega) \cap \mathbf{H}_0^1(\Omega), \mathbf{V}_n^0(\Omega)]_{1-r/2}, \quad \mathbf{V}^{-r}(\Omega) \stackrel{\text{def}}{=} [\mathbf{V}_0^r(\Omega)]',$$

and we recall that the orthogonal of $\mathbf{V}_n^0(\Omega)$ in $\mathbf{L}^2(\Omega)$ is given by:

$$[\mathbf{V}_n^0(\Omega)]^\perp = \{\nabla p \mid p \in H^1(\Omega)\}. \quad (2.1)$$

Standard arguments guarantee that $\mathbf{V}_0^r(\Omega) = \mathbf{V}_n^r(\Omega)$ if $r \in [0, 1/2)$, that $\mathbf{V}_0^r(\Omega) = \{y \in \mathbf{H}^r(\Omega) \cap \mathbf{V}_n^0(\Omega) \mid y = 0 \text{ on } \partial\Omega\}$ if $r \in (\frac{1}{2}, 2]$ and that $\mathbf{V}_0^{1/2}(\Omega) = \mathbf{V}_n^{1/2}(\Omega) \cap \mathbf{H}_{00}^{1/2}(\Omega)$. Next, we recall that the family $g_i, i = 1, \dots, N$ denotes a basis of the space X_N defined by (1.10), for $r \in [0, 2]$ we define

$$\widehat{\mathbf{V}}_n^r(\Omega) \stackrel{\text{def}}{=} \left\{ y \in \mathbf{V}_n^r(\Omega) \mid \int_{\Omega} y \cdot g_i dx = 0 \quad i = 1, \dots, N \right\},$$

and we recall that the following Poincaré-type inequality holds

$$\forall y \in \widehat{\mathbf{V}}_n^1(\Omega) \quad \int_{\Omega} |\text{curl } y(t)|^2 dx d\tau \geq c \int_{\Omega} (|\nabla y(t)|^2 + |y(t)|^2) dx, \quad (2.2)$$

where $c > 0$ is a positive constant depending only on Ω , see [19, Lemme 1.6].

Moreover, the orthogonal of $\widehat{\mathbf{V}}_n^0(\Omega)$ in $\mathbf{L}^2(\Omega)$ is given by (see [19]):

$$[\widehat{\mathbf{V}}_n^0(\Omega)]^\perp = \left\{ \nabla p + \sum_{i=1}^N \beta_i g_i \mid p \in H^1(\Omega), \beta_i \in \mathbb{R}, i = 1, \dots, N \right\}. \quad (2.3)$$

Finally, we set $Q \stackrel{\text{def}}{=} (0, T) \times \Omega$, $\Sigma \stackrel{\text{def}}{=} (0, T) \times \partial\Omega$, for $r \geq 0$ we define:

$$\begin{aligned} \mathbf{H}^{r, \frac{r}{2}}(Q) &\stackrel{\text{def}}{=} L^2(0, T; \mathbf{H}^r(\Omega)) \cap H^{\frac{r}{2}}(0, T; \mathbf{L}^2(\Omega)), \\ \mathbf{H}^{r, \frac{r}{2}}(\Sigma) &\stackrel{\text{def}}{=} L^2(0, T; \mathbf{H}^r(\partial\Omega)) \cap H^{\frac{r}{2}}(0, T; \mathbf{L}^2(\partial\Omega)), \end{aligned}$$

and for two Hilbert spaces H_1 and H_2 we set:

$$W(0, T; H_1, H_2) \stackrel{\text{def}}{=} L^2(0, T; H_1) \cap H^1(0, T; H_2).$$

3. Main steps of the proof of Theorem 1.1

The proof of Theorem 1.1 relies on a global Carleman inequality for the adjoint system (1.20). Let us introduce some usual weight functions that appear in Carleman inequalities for parabolic systems. Let $\omega \subset \mathcal{O}$ a nonempty open subset such that $\bar{\omega} \subset \mathcal{O}$ and let $\eta \in C^2(\bar{\Omega})$ such that

$$\begin{aligned} \eta(x) &= 0 \quad \forall x \in \partial\Omega, \\ \eta(x) &> 0 \quad \forall x \in \Omega, \\ |\nabla \eta(x)| &> 0 \quad \forall x \in \bar{\Omega} \setminus \omega. \end{aligned}$$

For the existence of such a function see for instance [24] or [42, Appendix III]. Thus, we introduce $e \in C^\infty([0, 1])$ such that $e(t) = t$ for $t \in (0, 1/4)$, $e(t) = 1 - t$ for $t \in (3/4, 1)$ and $e(t) \in [1/4, 1/2]$ for $t \in (1/4, 3/4)$, and we define $\ell(t) = Te(t/T)$. It is obvious to see that:

$$\ell \in C^\infty([0, T]) \quad \text{and} \quad \begin{cases} \ell(t) = t & \forall t \in (0, T/4), \\ \ell(t) \in [T/4, T/2] & \forall t \in (T/4, 3T/4), \\ \ell(t) = T - t & \forall t \in (3T/4, T). \end{cases} \quad (3.1)$$

Thus, for $k \geq 2$ and $\lambda > 1$ we define:

$$\begin{aligned}\alpha(t, x) &\stackrel{\text{def}}{=} \frac{e^{\lambda(\eta(x)+(k+1)\|\eta\|_\infty)} - e^{\lambda(k+2+\frac{1}{k})\|\eta\|_\infty}}{\ell(t)^k}, \\ \varphi(t, x) &\stackrel{\text{def}}{=} \frac{e^{\lambda(\eta(x)+(k+1)\|\eta\|_\infty)}}{\ell(t)^k},\end{aligned}\tag{3.2}$$

and

$$\begin{aligned}\widehat{\alpha}(t) &\stackrel{\text{def}}{=} \min_{x \in \overline{\Omega}} \alpha(t, x) = \frac{e^{\lambda(k+1)\|\eta\|_\infty} - e^{\lambda(k+2+\frac{1}{k})\|\eta\|_\infty}}{\ell(t)^k}, \\ \widehat{\varphi}(t) &\stackrel{\text{def}}{=} \min_{x \in \overline{\Omega}} \varphi(t, x) = \frac{e^{\lambda(k+1)\|\eta\|_\infty}}{\ell(t)^k}.\end{aligned}\tag{3.3}$$

The above weight functions satisfy

$$T^{-1}|\ell(t)| + |(\ell)_t(t)| + T|(\ell)_{tt}(t)| \leq c,\tag{3.4}$$

and for all $\lambda > 1$, $i = 1, 2, 3$ and $b \geq a$:

$$\lambda^{-2}|\partial_{x_i}^2 \varphi(t, x)| + \lambda^{-1}|\partial_{x_i} \varphi(t, x)| \leq c\varphi(t, x),\tag{3.5}$$

$$|\varphi^a(t, x)| \leq cT^{(b-a)k}\varphi^b(t, x),\tag{3.6}$$

$$|\varphi_t(t, x)| + |\alpha_t(t, x)| \leq c\varphi^{1+\frac{1}{k}}(t, x),\tag{3.7}$$

$$|\varphi_t(t, x)| + |\alpha_t(t, x)| \leq cT^{k-1}\varphi^2(t, x),\tag{3.8}$$

$$|\alpha_{tt}(t, x)| \leq cT^{2k-2}\varphi^3(t, x),\tag{3.9}$$

for some constant $c > 0$ which does not depend on T or λ .

The following theorem states a Carleman inequality for the adjoint system (1.20). Its proof is postponed to Section 4.

Theorem 3.1. *Assume $k \geq 4$. There exists $c_0 > 0$, $\lambda_0 > 0$ and $s_0 > 0$ such that for all $s \geq s_0$ and $\lambda \geq \lambda_0$ every solution of (1.20) satisfies:*

$$\begin{aligned}&\int_Q e^{2s\alpha} |\nabla(\text{curl } y)|^2 dx d\tau + s\lambda^2 \int_Q e^{2s\alpha} \varphi |\nabla y|^2 dx d\tau \\ &+ s^2 \lambda^2 \int_Q e^{2s\alpha} \varphi^2 |\text{curl } y|^2 dx d\tau + s^3 \lambda^4 \int_Q e^{2s\alpha} \varphi^3 |y|^2 dx d\tau \\ &+ \lambda \int_Q e^{2s\alpha} |\nabla(\text{curl } \rho)|^2 dx d\tau + s^2 \lambda^3 \int_Q e^{2s\alpha} \varphi^2 |\text{curl } \rho|^2 dx d\tau \\ &\leq c_0 \left(s \int_Q e^{2s\alpha} \varphi (|f_1|^2 + \lambda |f_2|^2) dx d\tau \right. \\ &\quad \left. + s^4 \lambda^4 \int_0^T \int_{\mathcal{O}} e^{2s\alpha} \varphi^4 (|y|^2 + \lambda |P_{\mathcal{O}} \rho|^2) dx d\tau \right).\end{aligned}\tag{3.10}$$

Next, we introduce $\tilde{\ell} \in C^\infty([0, T])$ such that $\tilde{\ell} \equiv \ell$ on $(\frac{T}{2}, T]$ and $\tilde{\ell} > 0$ on $[0, \frac{T}{2}]$ and we set

$$\widehat{\beta}(t) \stackrel{\text{def}}{=} \frac{e^{\lambda(k+1)\|\eta\|_\infty} - e^{\lambda(k+2+\frac{1}{k})\|\eta\|_\infty}}{\tilde{\ell}(t)^k}.$$

It is clear that for λ large enough we have:

$$\forall(t, x) \in \left[\frac{T}{2}, T \right] \times \bar{\Omega} \quad \alpha(t, x) < \frac{3}{4} \widehat{\beta}(t). \quad (3.11)$$

In the remaining part of this section we fix $k \geq 4$ and s, λ large enough so that (3.10) and (3.11) are satisfied.

The following theorem follows by combining the use of Theorem 6.11 in appendix and of Theorem 3.1 with (3.10), (3.11) and (2.2).

Theorem 3.2. *There exists $c > 0$ such that every solution of (1.20) satisfies:*

$$\begin{aligned} & \int_Q e^{2s\widehat{\beta}} (|\nabla y|^2 + |\nabla \rho|^2 + |y|^2 + |\rho|^2) dx d\tau + \|(y(0), \rho(0))\|_{\mathbf{V}_n^0(\Omega) \times \widehat{\mathbf{V}}_n^0(\Omega)} \\ & \leq c \left(\int_Q e^{s\frac{3\widehat{\beta}}{2}} (|f_1|^2 + |f_2|^2) dx d\tau + \int_0^T \int_{\mathcal{O}} e^{s\frac{3\widehat{\beta}}{2}} (|y|^2 + |P_{\mathcal{O}}\rho|^2) dx d\tau \right). \end{aligned} \quad (3.12)$$

The proof of the above theorem consists in combining the use of estimate (6.20) in appendix on $(\psi y, \psi \rho)$, where ψ is a smooth nonnegative cut-off function such that $\psi \equiv 1$ on $(0, T/2)$ and $\psi \equiv 0$ on $(3T/4, T)$, and the use of the Carleman inequality (3.10) together with (3.11) and (2.2). Since it is very classical, we omit it and we refer to [17, Lemma 1] or [28, Lemma 2] for a completely similar proofs.

The Theorem 3.2 permits to obtain a solution of the nonhomogeneous controlled system (1.19). We introduce the following weight spaces:

$$\begin{aligned} \mathcal{G} & \stackrel{\text{def}}{=} \{ (g_1, g_2) \in L^2(0, T; \mathbf{H}^{-1/2}(\Omega)) \times [\mathbf{H}_n^{1/2}(\Omega)]' \mid \\ & \quad (e^{-s\widehat{\beta}} g_1, e^{-s\widehat{\beta}} g_2) \in L^2(0, T; \mathbf{V}^{-1/2}(\Omega) \times [\mathbf{H}_n^{1/2}(\Omega)]' \}, \\ \mathcal{W} & \stackrel{\text{def}}{=} \{ (w, \theta) \in W(0, T; \mathbf{V}_0^{3/2}(\Omega) \times \widehat{\mathbf{V}}_n^{3/2}(\Omega), \mathbf{V}^{-1/2}(\Omega) \times [\widehat{\mathbf{V}}_n^{1/2}(\Omega)]' \mid \\ & \quad (e^{-s\frac{\widehat{\beta}}{2}} w, e^{-s\frac{\widehat{\beta}}{2}} \theta) \in W(0, T; \mathbf{V}_0^{3/2}(\Omega) \times \widehat{\mathbf{V}}_n^{3/2}(\Omega), \mathbf{V}^{-1/2}(\Omega) \times [\widehat{\mathbf{V}}_n^{1/2}(\Omega)]' \} \\ & \quad (e^{-s\frac{3\widehat{\beta}}{2}} w, e^{-s\frac{3\widehat{\beta}}{2}} \theta) \in \mathbf{L}^2(Q) \times \mathbf{L}^2(Q) \} \\ \mathcal{U} & \stackrel{\text{def}}{=} \{ (u_1, u_2) \in \mathbf{L}^2((0, T) \times \mathcal{O}) \times \mathbf{L}^2((0, T) \times \mathcal{O}) \mid \\ & \quad (e^{-s\frac{3\widehat{\beta}}{2}} u_1, e^{-s\frac{3\widehat{\beta}}{2}} u_2) \in \mathbf{L}^2((0, T) \times \mathcal{O}) \times \mathbf{L}^2((0, T) \times \mathcal{O}) \}, \end{aligned}$$

equipped with the norms

$$\begin{aligned} \|(g_1, g_2)\|_{\mathcal{G}} & \stackrel{\text{def}}{=} \|(e^{-s\widehat{\beta}} g_1, e^{-s\widehat{\beta}} g_2)\|_{L^2((0, T); \mathbf{H}^{-1/2}(\Omega)) \times [\mathbf{H}_n^{1/2}(\Omega)]'}, \\ \|(w, \theta)\|_{\mathcal{W}} & \stackrel{\text{def}}{=} \|(e^{-s\frac{\widehat{\beta}}{2}} w, e^{-s\frac{\widehat{\beta}}{2}} \theta)\|_{W(0, T; \mathbf{V}_0^{3/2}(\Omega) \times \widehat{\mathbf{V}}_n^{3/2}(\Omega), \mathbf{V}^{-1/2}(\Omega) \times [\widehat{\mathbf{V}}_n^{1/2}(\Omega)]' } \\ & \quad + \|(e^{-s\frac{3\widehat{\beta}}{2}} w, e^{-s\frac{3\widehat{\beta}}{2}} \theta)\|_{\mathbf{L}^2(Q)}, \\ \|(u_1, u_2)\|_{\mathcal{U}} & \stackrel{\text{def}}{=} \|(e^{-s\frac{3\widehat{\beta}}{2}} u_1, e^{-s\frac{3\widehat{\beta}}{2}} u_2)\|_{\mathbf{L}^2((0, T) \times \mathcal{O}) \times \mathbf{L}^2((0, T) \times \mathcal{O})}. \end{aligned}$$

We have the following null controllability result for the nonhomogeneous magnetohydrodynamic equations.

Theorem 3.3. *There exists $c_0 > 0$ such that for all $(w_0, \theta_0) \in \mathbf{V}_0^{1/2}(\Omega) \times \widehat{\mathbf{V}}_n^{1/2}(\Omega)$ and $(g_1, g_2) \in \mathcal{G}$ the control problem (1.19) admits a solution $(w, \theta, u_1, u_2) \in \mathcal{W} \times \mathcal{U}$ satisfying:*

$$\|(w, \theta)\|_{\mathcal{W}} + \|(u_1, u_2)\|_{\mathcal{U}} \leq c_0 (\|(g_1, g_2)\|_{\mathcal{G}} + \|(w_0, \theta_0)\|_{\mathbf{V}_0^{1/2}(\Omega) \times \widehat{\mathbf{V}}_n^{1/2}(\Omega)}).$$

Finally, the Theorem 1.1 is an obvious consequence of the following theorem which is obtained from Theorem 3.3 with a fixed-point argument.

Theorem 3.4. *There exist $c_0 > 0$ and $\varepsilon > 0$ such that for all $(w_0, \theta_0) \in \mathbf{V}_0^{1/2}(\Omega) \times \widehat{\mathbf{V}}_n^{1/2}(\Omega)$ such that*

$$\|(w_0, \theta_0)\|_{\mathbf{V}_0^{1/2}(\Omega) \times \widehat{\mathbf{V}}_n^{1/2}(\Omega)} \leq \varepsilon$$

system (1.17)-(1.18) admits a solution $(w, \theta, u_1, u_2) \in \mathcal{W} \times \mathcal{U}$ satisfying:

$$\|(w, \theta)\|_{\mathcal{W}} + \|(u_1, u_2)\|_{\mathcal{U}} \leq c_0 \|(w_0, \theta_0)\|_{\mathbf{V}_0^{1/2}(\Omega) \times \widehat{\mathbf{V}}_n^{1/2}(\Omega)}.$$

Moreover, if $(\tilde{w}, \tilde{\theta}) \in C(0, T; \mathbf{H}^{1/2}(\Omega) \times \mathbf{H}^{1/2}(\Omega)) \cap L^2(0, T; \mathbf{H}^{3/2}(\Omega) \times \mathbf{H}^{3/2}(\Omega))$ is another solution of (1.17) corresponding to (w_0, θ_0) and (u_1, u_2) then $(\tilde{w}, \tilde{\theta})$ and (w, θ) must coincide, i.e $(\tilde{w}, \tilde{\theta}) \equiv (w, \theta)$.

4. A Carleman inequality for the adjoint system

4.1. Main steps of the proof of Theorem 3.1

We set $\zeta \stackrel{\text{def}}{=} \text{curl} \rho$ and we apply the operator curl to the second equality in (1.20). By using the identities $(D^a \rho) \bar{B} = S(\bar{B}) \text{curl} \rho$ and $(D^a \rho) \bar{v} = S(\bar{v}) \text{curl} \rho$, where

$$S(z) \stackrel{\text{def}}{=} \begin{pmatrix} 0 & z_3 & -z_2 \\ -z_3 & 0 & z_1 \\ z_2 & -z_1 & 0 \end{pmatrix},$$

we get

$$\begin{cases} -y_t - \Delta y - (D^s y) \bar{v} + S(\bar{B}) \zeta + \nabla \pi = f_1 & \text{in } Q, \\ -\zeta_t - \Delta \zeta + \text{curl}((D^s y) \bar{B} - S(\bar{v}) \zeta) = \text{curl} f_2 & \text{in } Q, \\ \text{div } y = \text{div } \zeta = 0 & \text{in } Q, \\ y = \zeta \times n = 0 & \text{on } \Sigma. \end{cases} \quad (4.1)$$

Inequality (3.10) will be obtained by coupling the use of a Carleman inequality for Stokes equations and of a Carleman inequality for the curl of a Dynamo-type equations.

First, consider the adjoint Stokes equations with an homogeneous Dirichlet boundary condition:

$$\begin{cases} -\partial_t y - \Delta y + \nabla \pi = f & \text{in } Q, \\ \text{div } y = 0 & \text{in } Q, \\ y = 0 & \text{on } \Sigma. \end{cases} \quad (4.2)$$

The following theorem can be found in [36] (see also [35, 3]).

Theorem 4.1. *Assume $k \geq 4$. There exist $c_0 > 0$, $\lambda_0 > 0$ and $s_0 > 0$ such that for all $s \geq s_0$ and $\lambda \geq \lambda_0$ every solution of (4.2) satisfies*

$$\begin{aligned} & \int_Q e^{2s\alpha} |\nabla(\operatorname{curl} y)|^2 dx d\tau + s\lambda^2 \int_Q e^{2s\alpha} \varphi |\nabla y|^2 dx d\tau \\ & + s^2 \lambda^2 \int_Q e^{2s\alpha} \varphi^2 |\operatorname{curl} y|^2 dx d\tau + s^3 \lambda^4 \int_Q e^{2s\alpha} \varphi^3 |y|^2 dx d\tau \\ & \leq c_0 \left(s \int_Q e^{2s\alpha} \varphi |f|^2 dx d\tau + s^4 \lambda^4 \int_0^T \int_\Omega e^{2s\alpha} \varphi^4 |y|^2 dx d\tau \right). \end{aligned} \quad (4.3)$$

Next, consider the following equations:

$$\begin{cases} -\partial_t \zeta - \Delta \zeta & = \operatorname{curl} f & \text{in } Q, \\ \operatorname{div} \zeta & = 0 & \text{in } Q, \\ \zeta \times n & = 0 & \text{on } \Sigma. \end{cases} \quad (4.4)$$

To prove a Carleman inequality for system (4.4) we use the trick of [38]. For $m > 0$ we set $\xi = e^{m\eta} \zeta$. Since $\operatorname{div} \zeta = 0$ in Q we deduce that $\operatorname{div} \xi = m \nabla \eta \cdot \xi$ in Q and combining this with the fact that $\operatorname{div} \xi|_{\partial\Omega} = \operatorname{div}_\tau \xi_\tau + (\operatorname{div} n) \xi_n + \frac{\partial \xi_n}{\partial n}$ on $\partial\Omega$ (where div_τ is the surfacic divergence operator) from $\xi_\tau = 0$ we deduce that ξ satisfies

$$\frac{\partial \xi_n}{\partial n} + \left(\operatorname{div} n - m \frac{\partial \eta}{\partial n} \right) \xi_n = 0 \text{ on } \Sigma.$$

Thus, if we set

$$\gamma = \operatorname{div} n - m \frac{\partial \eta}{\partial n} \quad (4.5)$$

and

$$g_1 = -2m \nabla \xi \nabla \eta - (m e^{m\eta} \nabla \eta \times f + m^2 |\nabla \eta|^2 - m \Delta \eta) \xi, \quad g_2 = e^{m\eta} f, \quad (4.6)$$

then ξ satisfies the following vectorial heat equation with a Robin boundary condition on ξ_n and with a Dirichlet boundary condition on ξ_τ :

$$\begin{cases} -\partial_t \xi - \Delta \xi & = g_1 + \operatorname{curl}(g_2) & \text{in } Q, \\ \frac{\partial \xi_n}{\partial n} + \gamma \xi_n & = 0 & \text{on } \Sigma, \\ \xi_\tau & = 0 & \text{on } \Sigma. \end{cases} \quad (4.7)$$

The following Theorem states a Carleman inequality for system (4.7). Its proof is postponed to the Subsection 4.2.

Theorem 4.2. *Assume $k \geq 2$ and suppose that $\min_{\partial\Omega} \gamma(x) > 0$. There exist $c_0 > 0$, $\lambda_0 > 0$ and $s_0 > 0$ such that for all $s \geq s_0$ and $\lambda \geq \lambda_0$ every solution of (4.7) satisfies*

$$\begin{aligned} & s\lambda^2 \int_Q e^{2s\alpha} \varphi |\nabla \xi|^2 dx d\tau + s^3 \lambda^4 \int_Q e^{2s\alpha} \varphi^3 |\xi|^2 dx d\tau \\ & \leq c_0 \left(\int_Q e^{2s\alpha} |g_1|^2 dx d\tau + s^2 \lambda^2 \int_Q e^{2s\alpha} \varphi^2 |g_2|^2 dx d\tau \right. \\ & \quad \left. + s^3 \lambda^4 \int_0^T \int_\omega e^{2s\alpha} \varphi^3 |\xi|^2 dx d\tau \right). \end{aligned} \quad (4.8)$$

To deduce a Carleman inequality for system (4.4) from the above theorem we choose m large enough so that

$$-\frac{m}{2} \frac{\partial \eta}{\partial n} \leq \gamma \leq -2m \frac{\partial \eta}{\partial n}. \quad (4.9)$$

Theorem 4.3. *Assume $k \geq 2$. There exist $c_0 > 0$, $\lambda_0 > 0$ and $s_0 > 0$ such that for all $s \geq s_0$ and $\lambda \geq \lambda_0$ every solution of (4.4) satisfies*

$$\begin{aligned} & \lambda \int_Q e^{2s\alpha} |\nabla \zeta|^2 dx d\tau + s^2 \lambda^3 \int_Q e^{2s\alpha} \varphi^2 |\zeta|^2 dx d\tau \\ & \leq c_0 \left(s \lambda \int_Q \varphi e^{2s\alpha} |f|^2 dx d\tau + s^2 \lambda^3 \int_0^T \int_\omega e^{2s\alpha} \varphi^2 |\zeta|^2 dx d\tau \right). \end{aligned} \quad (4.10)$$

Proof. By applying (4.8) to (4.7)-(4.6) and recalling (4.9) we obtain

$$\begin{aligned} & s \lambda^2 \int_Q e^{2s\alpha} \varphi |\nabla \xi|^2 dx d\tau + s^3 \lambda^4 \int_Q e^{2s\alpha} \varphi^3 |\xi|^2 dx d\tau \\ & \leq c_0 \left(\int_Q e^{2s\alpha} (m^2 e^{2m\eta} |f|^2 + m^4 |\xi|^2 + m^2 |\nabla \xi|^2) dx d\tau \right. \\ & \quad \left. + s^2 \lambda^2 \int_Q e^{2s\alpha} \varphi^2 e^{2m\eta} |f|^2 dx d\tau + s^3 \lambda^4 \int_0^T \int_\omega e^{2s\alpha} \varphi^3 |\xi|^2 dx d\tau \right) \end{aligned}$$

and we conclude by choosing s large enough and recalling that $\zeta = e^{-m\eta} \xi$. \square

Remark 4.4. The strategy of proof consisting in reducing (4.4) to system (4.7) is specific to the three dimensional case and proving Theorem 4.2 is useless in the two dimensional case. Indeed, in the two dimensional counterpart of (1.20) the boundary condition on $\text{curl} \rho$ is simply $\text{curl} \rho = 0$ on Σ and (4.4) must be rewritten as

$$\begin{cases} -\partial_t \zeta - \Delta \zeta & = \text{curl} f & \text{in } Q, \\ \text{div} \zeta & = 0 & \text{in } Q, \\ \zeta & = 0 & \text{on } \Sigma. \end{cases} \quad (4.11)$$

Then inequality (4.10) follows from a direct application of the Carleman inequality for parabolic equations with source term in $H^{-1}(\Omega)$ proved in [33], see also [15, Lem. 2.1].

We are now in position to prove Theorem 3.1.

Proof of Theorem 3.1. If we apply (4.3) and (4.10) to the first and the second equality of (4.1) respectively and if we choose $\lambda \geq c(\|\bar{v}\|_{\mathbf{L}^\infty(Q)} + \|\bar{B}\|_{\mathbf{L}^\infty(Q)})$

and $c > 0$ large enough we obtain

$$\begin{aligned}
& \int_Q e^{2s\alpha} |\nabla(\operatorname{curl} y)|^2 dx d\tau + s\lambda^2 \int_Q e^{2s\alpha} \varphi |\nabla y|^2 dx d\tau \\
& + s^2 \lambda^2 \int_Q e^{2s\alpha} \varphi^2 |\operatorname{curl} y|^2 dx d\tau + s^3 \lambda^4 \int_Q e^{2s\alpha} \varphi^3 |y|^2 dx d\tau \\
& + \lambda \int_Q e^{2s\alpha} |\nabla(\operatorname{curl} \rho)|^2 dx d\tau + s^2 \lambda^3 \int_Q e^{2s\alpha} \varphi^2 |\operatorname{curl} \rho|^2 dx d\tau \\
& \leq C \left(s \int_Q e^{2s\alpha} \varphi (|f_1|^2 + \lambda |f_2|^2) dx d\tau + s^4 \lambda^4 \int_0^T \int_{\mathcal{O}} e^{2s\alpha} \varphi^3 |y|^2 dx d\tau \right. \\
& \quad \left. + s^2 \lambda^3 \int_0^T \int_{\omega} e^{2s\alpha} \varphi^2 |\operatorname{curl} \rho|^2 dx d\tau \right) \tag{4.12}
\end{aligned}$$

for $C > 0$ independent on s, λ . Next, we use a classical localization argument to obtain the second local term at the right of inequality (3.10). We recall that $\bar{\omega} \subset \mathcal{O}$ and let $\vartheta \in C_c^\infty(\mathcal{O})$, $0 \leq \vartheta \leq 1$ and $\vartheta \equiv 1$ in ω . First, we have:

$$s^2 \int_0^T \int_{\omega} e^{2s\alpha} \varphi^2 |\operatorname{curl} \rho|^2 dx d\tau \leq s^2 \int_0^T \int_{\mathcal{O}} \vartheta e^{2s\alpha} \varphi^2 |\operatorname{curl} \rho|^2 dx d\tau.$$

We recall that $P_{\mathcal{O}}\rho = \rho - \nabla\chi$ for some $\chi \in H^1(\Omega)$ and it implies $\operatorname{curl} \rho = \operatorname{curl} P_{\mathcal{O}}\rho$. Then with the following computations:

$$\begin{aligned}
s^2 \int_0^T \int_{\mathcal{O}} \vartheta e^{2s\alpha} \varphi^2 |\operatorname{curl} \rho|^2 dx d\tau &= s^2 \int_0^T \int_{\mathcal{O}} \vartheta e^{2s\alpha} \varphi^2 \operatorname{curl} \rho \cdot \operatorname{curl} P_{\mathcal{O}}\rho dx d\tau \\
&= s^2 \int_0^T \int_{\mathcal{O}} \vartheta e^{2s\alpha} \varphi^2 \operatorname{curl}(\operatorname{curl} \rho) \cdot P_{\mathcal{O}}\rho dx d\tau \\
&+ s^2 \int_0^T \int_{\mathcal{O}} e^{2s\alpha} \varphi^2 (\nabla\vartheta \times (\operatorname{curl} \rho)) \cdot P_{\mathcal{O}}\rho dx d\tau \\
&+ 2\lambda s^2 \int_0^T \int_{\mathcal{O}} \vartheta e^{2s\alpha} \varphi^2 (\nabla\eta \times (\operatorname{curl} \rho)) \cdot P_{\mathcal{O}}\rho dx d\tau \\
&+ 2\lambda s^3 \int_0^T \int_{\mathcal{O}} \vartheta e^{2s\alpha} \varphi^3 (\nabla\eta \times (\operatorname{curl} \rho)) \cdot P_{\mathcal{O}}\rho dx d\tau,
\end{aligned}$$

we obtain with (3.6) and $\epsilon > 0$:

$$\begin{aligned}
s^2 \lambda^3 \int_0^T \int_{\omega} e^{2s\alpha} \varphi |\operatorname{curl} \rho|^2 dx d\tau &\leq \epsilon \lambda \int_Q e^{2s\alpha} |\nabla(\operatorname{curl} \rho)|^2 dx d\tau \\
&+ \epsilon s^2 \lambda^3 \int_Q e^{2s\alpha} \varphi^2 |\operatorname{curl} \rho|^2 dx d\tau + \frac{C}{\epsilon} s^4 \lambda^5 \int_0^T \int_{\mathcal{O}} e^{2s\alpha} \varphi^4 |P_{\mathcal{O}}\rho|^2 dx d\tau.
\end{aligned}$$

Then with (4.12) and $\epsilon > 0$ small enough we get (3.10). \square

4.2. A Carleman inequality for a vectorial heat equation with mixed Robin and Dirichlet boundary conditions

The goal of this subsection is to prove Theorem 4.2. In the whole subsection C denotes a generic positive constant that may change from line to line, which

depends on the geometry, but which is independent on T , s , λ . Moreover, we introduce the following boundary integral:

$$\begin{aligned} I_\Sigma(s, \lambda, \psi) &\stackrel{\text{def}}{=} -2s^3\lambda^3 \int_\Sigma |\nabla\eta|^2 \widehat{\varphi}^3 \frac{\partial\eta}{\partial n} |\psi|^2 d\Sigma + 2s^2\lambda \int_\Sigma \widehat{\alpha}_t \widehat{\varphi} \frac{\partial\eta}{\partial n} |\psi|^2 d\Sigma \\ &\quad - 4s\lambda \int_\Sigma \widehat{\varphi} \frac{\partial\eta}{\partial n} \left| \frac{\partial\psi}{\partial n} \right|^2 d\Sigma + 2s\lambda \int_\Sigma \widehat{\varphi} \frac{\partial\eta}{\partial n} |\nabla\psi|^2 d\Sigma \\ &\quad - 4s\lambda^2 \int_\Sigma \left| \frac{\partial\eta}{\partial n} \right|^2 \widehat{\varphi} \frac{\partial\psi}{\partial n} \psi d\Sigma + 2 \int_\Sigma \frac{\partial\psi}{\partial n} \psi_t d\Sigma. \end{aligned}$$

We will need the following Carleman inequality for the heat equation with a nonhomogeneous boundary condition.

Lemma 4.5. *There exist $\lambda_0 > 0$, $c_0 > 0$ and $c_1 > 0$ independent on T and such that for all $\lambda \geq \lambda_0$ and $s \geq c_1(T^k + T^{k-1})$ the following inequality holds for all $q \in C^2(\overline{Q})$:*

$$\begin{aligned} &s^{-1} \int_Q e^{2s\alpha} \varphi^{-1} (|\partial_t q|^2 + |\Delta q|^2) dx d\tau + s\lambda^2 \int_Q e^{2s\alpha} \varphi |\nabla q|^2 dx d\tau \\ &+ s^3 \lambda^4 \int_Q e^{2s\alpha} \varphi^3 |q|^2 dx d\tau + I_\Sigma(s, \lambda, e^{s\alpha} q) \\ &\leq c_0 \left(\int_Q e^{2s\alpha} |\partial_t q + \Delta q|^2 dx d\tau + s^3 \lambda^4 \int_0^T \int_\omega e^{2s\alpha} \varphi^3 |q|^2 dx d\tau \right). \end{aligned} \quad (4.13)$$

For the proof of Lemma 4.5 we refer to the Appendix of [14].

Next, to get rid of the boundary terms in (4.13) corresponding to powers of λ with odd exponents, we need another Carleman inequality involving weight functions where η is replaced by $-\eta$. For $k \geq 2$ and $\lambda > 1$ we define:

$$\begin{aligned} \tilde{\alpha}(t, x) &= \frac{e^{\lambda(-\eta(x)+(k+1)\|\eta\|_\infty)} - e^{\lambda(k+2+\frac{1}{k})\|\eta\|_\infty}}{\ell(t)^k}, \\ \tilde{\varphi}(t, x) &= \frac{e^{\lambda(-\eta(x)+(k+1)\|\eta\|_\infty)}}{\ell(t)^k}. \end{aligned} \quad (4.14)$$

We verify that $\tilde{\varphi}$ satisfies (3.5)-(3.6) and $\tilde{\alpha}$ satisfies

$$\begin{aligned} |\tilde{\alpha}_t(t, x)| &\leq C e^{\lambda(1+\frac{1}{k})\|\eta\|_\infty} \tilde{\varphi}^{1+\frac{1}{k}}(t, x), \\ |\tilde{\alpha}_{tt}(t, x)| &\leq C e^{\lambda\frac{1}{k}\|\eta\|_\infty} \tilde{\varphi}^{1+\frac{2}{k}}(t, x), \\ |\tilde{\alpha}_t(t, x)| &\leq C e^{\lambda(1+\frac{1}{k})\|\eta\|_\infty} T^{k-1} \tilde{\varphi}^2(t, x), \\ |\tilde{\alpha}_{tt}(t, x)| &\leq C e^{\lambda\frac{1}{k}\|\eta\|_\infty} T^{2k-2} \tilde{\varphi}^3(t, x). \end{aligned}$$

Note that we have

$$\tilde{\alpha}(t, x) = \alpha(t, x) = \widehat{\alpha}(t) \quad \text{and} \quad \tilde{\varphi}(t, x) = \varphi(t, x) = \widehat{\alpha}(t) \quad \forall x \in \partial\Omega. \quad (4.15)$$

The proof of the following lemma can also be found in the Appendix of [14].

Lemma 4.6. *There exist $\lambda_0 > 0$, $c_0 > 0$ and $c_1 > 0$ independent on T and such that for all $\lambda \geq \lambda_0$ and $s \geq c_1 e^{\lambda(1+\frac{1}{k})\|\eta\|_\infty}(T^k + T^{k-1})$ the following inequality holds for all $q \in C^2(\bar{Q})$:*

$$\begin{aligned}
& s^{-1} \int_Q e^{2s\tilde{\alpha}} \tilde{\varphi}^{-1} (|\partial_t q|^2 + |\Delta q|^2) dx d\tau + s\lambda^2 \int_Q e^{2s\tilde{\alpha}} \tilde{\varphi} |\nabla q|^2 dx d\tau \\
& + s^3 \lambda^4 \int_Q e^{2s\tilde{\alpha}} \tilde{\varphi}^3 |q|^2 dx d\tau + I_\Sigma(s, -\lambda, e^{s\tilde{\alpha}} q) \\
& \leq c_0 \left(\int_Q e^{2s\tilde{\alpha}} |\partial_t q + \Delta q|^2 dx d\tau + s^3 \lambda^4 \int_0^T \int_\omega e^{2s\tilde{\alpha}} \tilde{\varphi}^3 |q|^2 dx d\tau \right).
\end{aligned} \tag{4.16}$$

By combining the two previous lemmas we get the following theorem.

Theorem 4.7. *There exist $\lambda_0 > 0$, $c_0 > 0$ and $c_1 > 0$ independent on T and such that for all $\lambda \geq \lambda_0$ and $s \geq c_1 e^{\lambda(1+\frac{1}{k})\|\eta\|_\infty}(T^k + T^{k-1})$ the following inequality holds for all $q \in C^2(\bar{Q})$:*

$$\begin{aligned}
& s^{-1} \int_Q e^{2s\alpha} \varphi^{-1} (|\partial_t q|^2 + |\Delta q|^2) dx d\tau \\
& + s\lambda^2 \int_Q e^{2s\alpha} \varphi |\nabla q|^2 dx d\tau + s^3 \lambda^4 \int_Q e^{2s\alpha} \varphi^3 |q|^2 dx d\tau \\
& \leq c_0 \left(\int_Q e^{2s\alpha} |\partial_t q + \Delta q|^2 dx d\tau + s^3 \lambda^4 \int_0^T \int_\omega e^{2s\alpha} \varphi^3 |q|^2 dx d\tau \right) \\
& + 8s\lambda^2 \int_\Sigma e^{2s\hat{\alpha}} \hat{\varphi} \left| \frac{\partial \eta}{\partial n} \right|^2 (1 + s\hat{\varphi}) \frac{\partial q}{\partial n} q d\Sigma - 4 \int_\Sigma e^{s\hat{\alpha}} \frac{\partial q}{\partial n} (e^{s\hat{\alpha}} q)_t d\Sigma.
\end{aligned}$$

Proof. We add (4.13) and (4.16) and since $\tilde{\alpha} \leq \alpha$ and $\tilde{\varphi} \leq \varphi$ we first get

$$\begin{aligned}
& s^{-1} \int_Q e^{2s\alpha} \varphi^{-1} (|\partial_t q|^2 + |\Delta q|^2) dx d\tau + s\lambda^2 \int_Q e^{2s\alpha} \varphi |\nabla q|^2 dx d\tau \\
& + s^3 \lambda^4 \int_Q e^{2s\alpha} \varphi^3 |q|^2 dx d\tau + I_\Sigma(s, \lambda, e^{s\alpha} q) + I_\Sigma(s, -\lambda, e^{s\tilde{\alpha}} q) \\
& \leq C \left(\int_Q e^{2s\alpha} |\partial_t q + \Delta q|^2 dx d\tau + s^3 \lambda^4 \int_0^T \int_\omega e^{2s\alpha} \varphi^3 |q|^2 dx d\tau \right),
\end{aligned}$$

for $\lambda \geq \lambda_0$ and $s \geq C e^{\lambda(1+\frac{1}{k})\|\eta\|_\infty}(T^k + T^{k-1})$. Thus, from (4.15) we deduce that $\nabla(e^{s\alpha} q) = e^{s\hat{\alpha}}(s\lambda\hat{\varphi}\nabla\eta q + \nabla q)$, $\nabla(e^{s\tilde{\alpha}} q) = e^{s\hat{\alpha}}(-s\lambda\hat{\varphi}\nabla\eta q + \nabla q)$ and that

$(e^{s\alpha}q)_t = (e^{s\hat{\alpha}}q)_t = (e^{s\hat{\alpha}}q)_t$ on $\partial\Omega$, and we conclude with the calculations:

$$\begin{aligned}
& I_{\Sigma}(s, \lambda, e^{s\alpha}q) + I_{\Sigma}(s, -\lambda, e^{s\hat{\alpha}}q) \\
&= -4s\lambda \int_{\Sigma} e^{2s\hat{\alpha}} \hat{\varphi} \frac{\partial \eta}{\partial n} \left(\left| \frac{\partial q}{\partial n} + s\lambda \hat{\varphi} \frac{\partial \eta}{\partial n} q \right|^2 - \left| \frac{\partial q}{\partial n} - s\lambda \hat{\varphi} \frac{\partial \eta}{\partial n} q \right|^2 \right) d\Sigma \\
&+ 2s\lambda \int_{\Sigma} e^{2s\hat{\alpha}} \hat{\varphi} \frac{\partial \eta}{\partial n} (|\nabla q + s\lambda \hat{\varphi} \nabla \eta q|^2 - |\nabla q - s\lambda \hat{\varphi} \nabla \eta q|^2) d\Sigma \\
&- 4s\lambda^2 \int_{\Sigma} e^{2s\hat{\alpha}} \left| \frac{\partial \eta}{\partial n} \right|^2 \hat{\varphi} \left(\frac{\partial q}{\partial n} + s\lambda \hat{\varphi} \frac{\partial \eta}{\partial n} q \right) q d\Sigma \\
&- 4s\lambda^2 \int_{\Sigma} e^{2s\hat{\alpha}} \left| \frac{\partial \eta}{\partial n} \right|^2 \hat{\varphi} \left(\frac{\partial q}{\partial n} - s\lambda \hat{\varphi} \frac{\partial \eta}{\partial n} q \right) q d\Sigma \\
&+ \int_{\Sigma} e^{s\hat{\alpha}} \left(\frac{\partial q}{\partial n} + s\lambda \hat{\varphi} \frac{\partial \eta}{\partial n} q \right) (e^{s\hat{\alpha}}q)_t d\Sigma \\
&+ 2 \int_{\Sigma} e^{s\hat{\alpha}} \left(\frac{\partial q}{\partial n} - s\lambda \hat{\varphi} \frac{\partial \eta}{\partial n} q \right) (e^{s\hat{\alpha}}q)_t d\Sigma \\
&= -8s^2\lambda^2 \int_{\Sigma} e^{2s\hat{\alpha}} \hat{\varphi}^2 \left| \frac{\partial \eta}{\partial n} \right|^2 \frac{\partial q}{\partial n} q d\Sigma - 8s\lambda^2 \int_{\Sigma} e^{2s\hat{\alpha}} \hat{\varphi} \left| \frac{\partial \eta}{\partial n} \right|^2 \frac{\partial q}{\partial n} q d\Sigma \\
&+ 4 \int_{\Sigma} e^{s\hat{\alpha}} \frac{\partial q}{\partial n} (e^{s\hat{\alpha}}q)_t d\Sigma.
\end{aligned}$$

□

A consequence of Theorem 4.7 is the following Carleman inequality for a vectorial heat equation with mixed Robin and Dirichlet boundary conditions.

Corollary 4.8. *There exist $\lambda_0 > 0$, $c_0 > 0$ and $c_1 > 0$ independent on T and such that for all $\lambda \geq \lambda_0$ and $s \geq c_1 e^{\lambda(1+\frac{1}{k})\|\eta\|_{\infty}}(T^k + T^{k-1})$ the following inequality holds for all $\xi \in \mathbf{C}^2(\overline{Q})$ such that $\xi_{\tau} = 0$ and $\frac{\partial \xi_n}{\partial n} + \gamma \xi_n = 0$ on Σ :*

$$\begin{aligned}
& s^{-1} \int_Q e^{2s\alpha} \varphi^{-1} (|\partial_t \xi|^2 + |\Delta \xi|^2) dx d\tau + s\lambda^2 \int_Q e^{2s\alpha} \varphi |\nabla \xi|^2 dx d\tau \\
&+ s^3 \lambda^4 \int_Q e^{2s\alpha} \varphi^3 |\xi|^2 dx d\tau + s^2 \lambda^2 \int_{\Sigma} e^{2s\hat{\alpha}} \hat{\varphi}^2 \gamma |\xi_n|^2 d\Sigma \\
&\leq c_0 \left(\int_Q e^{2s\alpha} |\partial_t \xi + \Delta \xi|^2 dx d\tau + s^3 \lambda^4 \int_0^T \int_{\omega} e^{2s\alpha} \varphi^3 |\xi|^2 dx d\tau \right).
\end{aligned}$$

Proof. By applying Theorem 4.7 to each component of ξ we obtain:

$$\begin{aligned}
& s^{-1} \int_Q e^{2s\alpha} \varphi^{-1} (|\partial_t \xi|^2 + |\Delta \xi|^2) dx d\tau \\
& + s\lambda^2 \int_Q e^{2s\alpha} \varphi |\nabla \xi|^2 dx d\tau + s^3 \lambda^4 \int_Q e^{2s\alpha} \varphi^3 |\xi|^2 dx d\tau \\
& \leq C \left(\int_Q e^{2s\alpha} |\partial_t \xi + \Delta \xi|^2 dx d\tau + s^3 \lambda^4 \int_0^T \int_\omega e^{2s\alpha} \varphi^3 |\xi|^2 dx d\tau \right) \\
& + 4s\lambda^2 \int_\Sigma e^{2s\hat{\alpha}} \hat{\varphi} \left| \frac{\partial \eta}{\partial n} \right|^2 (1 + s\hat{\varphi}) \frac{\partial \xi}{\partial n} \cdot \xi d\Sigma - 2 \int_\Sigma e^{s\hat{\alpha}} \frac{\partial \xi}{\partial n} \cdot (e^{s\hat{\alpha}} \xi)_t d\Sigma,
\end{aligned} \tag{4.17}$$

and the conclusion follows from the fact that $\xi_\tau = 0$ and $\frac{\partial \xi_n}{\partial n} + \gamma \xi_n = 0$ yield

$$\int_\Sigma e^{s\hat{\alpha}} \frac{\partial \xi}{\partial n} \cdot (e^{s\hat{\alpha}} \xi)_t d\Sigma = -\frac{1}{2} \int_\Sigma \gamma ((e^{s\hat{\alpha}} \xi_n)_t)^2 d\Sigma = 0,$$

and from the fact that $\frac{\partial \xi}{\partial n} \cdot \xi = -\gamma |\xi_n|^2$ yields

$$4s\lambda^2 \int_\Sigma e^{2s\hat{\alpha}} \hat{\varphi} \left| \frac{\partial \eta}{\partial n} \right|^2 (1 + s\hat{\varphi}) \frac{\partial \xi}{\partial n} \cdot \xi d\Sigma \leq -Cs\lambda^2 \int_\Sigma e^{2s\hat{\alpha}} \hat{\varphi} (1 + s\hat{\varphi}) \gamma |\xi_n|^2 d\Sigma.$$

□

Finally, by duality we obtain a Carleman inequality for (4.7).

Theorem 4.9. *There exist $\lambda_0 > 0$, $c_0 > 0$ and $c_1 > 0$ independent on T and such that for all $\lambda \geq \lambda_0$ and $s \geq c_1 e^{\lambda(1+\frac{1}{k})\|\eta\|_\infty} (T^k + T^{k-1})$ every solution of (4.7) satisfies*

$$\begin{aligned}
& s\lambda^2 \int_Q e^{2s\alpha} \varphi |\nabla \xi|^2 dx d\tau + s^3 \lambda^4 \int_Q e^{2s\alpha} \varphi^3 |\xi|^2 dx d\tau \\
& \leq c_0 \left(\int_Q e^{2s\alpha} |g_1|^2 dx d\tau + s^2 \lambda^2 \int_Q e^{2s\alpha} \varphi^2 |g_2|^2 dx d\tau \right. \\
& \quad \left. + s^3 \lambda^4 \int_0^T \int_\omega e^{2s\alpha} \varphi^3 |\xi|^2 dx d\tau \right).
\end{aligned} \tag{4.18}$$

Proof. The proof follows from a now standard duality argument.

First, we construct a solution of

$$\begin{cases} \partial_t z - \Delta z & = s^3 \lambda^4 e^{2s\alpha} \varphi^3 \xi + \mathbf{1}_\omega u & \text{in } Q, \\ \frac{\partial z_n}{\partial n} + \gamma z_n & = 0 & \text{on } \Sigma, \\ z_\tau & = 0 & \text{on } \Sigma, \\ z(0) = z(T) & = 0 & \text{on } \Omega, \end{cases} \tag{4.19}$$

satisfying for $s \geq C(T^k + T^{k-1})$ and $\lambda > \lambda_0$, and $C > 0$, $\lambda_0 > 0$ large enough:

$$\begin{aligned}
& \int_Q e^{-2s\alpha} |z|^2 dx d\tau + s^{-3} \lambda^{-4} \int_0^T \int_\omega e^{-2s\alpha} \varphi^{-3} |u|^2 dx d\tau \\
& + s^{-2} \lambda^{-2} \int_Q e^{-2s\alpha} \varphi^{-2} |\nabla z|^2 dx d\tau \leq Cs^3 \lambda^4 \int_Q e^{2s\alpha} \varphi^3 |\xi|^2 dx d\tau.
\end{aligned} \tag{4.20}$$

For that we set

$$\mathcal{P}_0 \stackrel{\text{def}}{=} \left\{ w \in \mathbf{C}^2(\bar{Q}) \mid \frac{\partial w_n}{\partial n} + \gamma w_n = 0 \text{ and } w_\tau = 0 \text{ on } \Sigma \right\},$$

and we consider the bilinear form $b(\cdot, \cdot) : \mathcal{P}_0 \times \mathcal{P}_0 \rightarrow \mathbb{R}$ defined by

$$\begin{aligned} b(w, w') &= \int_Q e^{2s\alpha} (\partial_t w + \Delta w) \cdot (\partial_t w' + \Delta w') dx d\tau \\ &\quad + s^3 \lambda^4 \int_0^T \int_\omega e^{2s\alpha} \varphi^3 w \cdot w' dx d\tau, \end{aligned}$$

as well as the following linear form on \mathcal{P}_0 :

$$l(w') = s^3 \lambda^4 \int_Q e^{2s\alpha} \varphi^3 \xi \cdot w' dx d\tau.$$

The Corollary 4.8 guarantees that the semi-norm induced by $b(\cdot, \cdot)$ is a norm. If \mathcal{P} is the completion of \mathcal{P}_0 for this norm, then Corollary 4.8 guarantees the continuity of l on \mathcal{P} , and from the Lax-Milgram Lemma we obtain a unique solution $w \in \mathcal{P}$ to

$$b(w, w') = l(w'), \quad \forall w' \in \mathcal{P}. \quad (4.21)$$

Moreover, w satisfies

$$\begin{aligned} \int_Q e^{2s\alpha} |\partial_t w + \Delta w|^2 dx d\tau + s^3 \lambda^4 \int_0^T \int_\omega e^{2s\alpha} \varphi^3 |w|^2 dx d\tau \\ \leq C s^3 \lambda^4 \int_Q e^{2s\alpha} \varphi^3 |\xi|^2 dx d\tau. \end{aligned} \quad (4.22)$$

If we set $z = e^{2s\alpha} (\partial_t w + \Delta w)$ and $u = -s^3 \lambda^4 e^{2s\alpha} \varphi^3 w$ then (4.21) becomes:

$$\int_Q z \cdot (\partial_t w' + \Delta w') dx d\tau = s^3 \lambda^4 \int_Q e^{2s\alpha} \varphi^3 \xi \cdot w' dx d\tau + \int_Q \mathbf{1}_\omega u \cdot w' dx d\tau$$

which is the weak transposition formulation of (4.19), and (4.22) becomes

$$\begin{aligned} \int_Q e^{-2s\alpha} |z|^2 dx d\tau + s^{-3} \lambda^{-4} \int_0^T \int_\omega e^{-2s\alpha} \varphi^{-3} |u|^2 dx d\tau \leq \\ C s^3 \lambda^4 \int_Q e^{2s\alpha} \varphi^3 |\xi|^2 dx d\tau. \end{aligned} \quad (4.23)$$

Let us now give an estimate of ∇z . For $\epsilon > 0$ we introduce

$$\begin{aligned} \alpha_\epsilon(t, x) &\stackrel{\text{def}}{=} \frac{e^{\lambda(\eta(x) + (k+1)\|\eta\|_\infty)} - e^{\lambda(k+2+\frac{1}{k})\|\eta\|_\infty}}{(\ell(t) + \epsilon)^k}, \\ \varphi_\epsilon(t, x) &\stackrel{\text{def}}{=} \frac{e^{\lambda(\eta(x) + (k+1)\|\eta\|_\infty)}}{(\ell(t) + \epsilon)^k}. \end{aligned}$$

Thus, by multiplying the first equality in (4.19) by $s^{-2}\lambda^{-2}e^{-2s\alpha_\epsilon}\varphi_\epsilon^{-2}z$ and integrating by parts ($\alpha_\epsilon, \varphi_\epsilon$ do not blow up as $t \rightarrow 0$ or T) we obtain:

$$\begin{aligned} & s^{-2}\lambda^{-2} \int_Q e^{-2s\alpha_\epsilon}\varphi_\epsilon^{-2}|\nabla z|^2 dx d\tau + s^{-2}\lambda^{-2} \int_\Sigma e^{-2s\hat{\alpha}_\epsilon}\widehat{\varphi}_\epsilon^{-2}\gamma|z_n|^2 d\Sigma = \\ & s^{-2}\lambda^{-2} \int_0^T \int_\omega e^{-2s\alpha_\epsilon}\varphi_\epsilon^{-2}u \cdot z dx d\tau - s^{-2}\lambda^{-2} \int_Q (z^t \nabla(e^{-2s\alpha_\epsilon}\varphi_\epsilon^{-2})) : \nabla z dx d\tau \\ & + \frac{1}{2}s^{-2}\lambda^{-2} \int_Q (e^{-2s\alpha_\epsilon}\varphi_\epsilon^{-2})_t |z|^2 dx d\tau + s\lambda^2 \int_Q \varphi_\epsilon \xi \cdot z dx d\tau. \end{aligned}$$

Moreover, for $s \geq C(T^k + T^{k-1})$ and $C > 0$ large enough we verify that have

$$\begin{aligned} s^{-2}\lambda^{-2} |(e^{-2s\alpha_\epsilon}\varphi_\epsilon^{-2})_t| & \leq C\lambda^{-2}e^{-2s\alpha_\epsilon}, \\ s^{-2}\lambda^{-2} |\nabla(e^{-2s\alpha_\epsilon}\varphi_\epsilon^{-2})| & \leq Cs^{-1}\lambda^{-1}e^{-2s\alpha_\epsilon}\varphi_\epsilon^{-1}. \end{aligned}$$

Then combining this with the above equality we deduce that:

$$\begin{aligned} & s^{-2}\lambda^{-2} \int_Q e^{-2s\alpha_\epsilon}\varphi_\epsilon^{-2}|\nabla z|^2 dx d\tau \leq C \left(\int_Q e^{-2s\alpha_\epsilon}|z|^2 dx d\tau \right. \\ & \left. + s^{-3}\lambda^{-4} \int_0^T \int_\omega e^{-2s\alpha_\epsilon}\varphi_\epsilon^{-3}|u|^2 dx d\tau + s^3\lambda^4 \int_Q e^{2s\alpha_\epsilon}\varphi_\epsilon^3|\xi|^2 dx d\tau \right), \end{aligned}$$

for a constant $C > 0$ independent on ϵ . Then we can pass to the limit $\epsilon \rightarrow 0$ and with (4.23) it yields (4.20).

We are now in position to deduce the estimate (4.18) for ξ by duality. By multiplying the first equation in (4.7) by z and integrate by parts we get

$$\begin{aligned} s^3\lambda^4 \int_Q e^{2s\alpha}\varphi^3|\xi|^2 dx d\tau & = \int_Q g_1 \cdot z dx d\tau \\ & + \int_Q g_2 \cdot \text{curl} z dx d\tau - \int_0^T \int_\omega u \cdot \xi dx d\tau. \end{aligned} \tag{4.24}$$

Then the above equality with (4.20) yields:

$$\begin{aligned} s^3\lambda^4 \int_Q e^{2s\alpha}\varphi^3|\xi|^2 dx d\tau & \leq C \left(\int_Q e^{2s\alpha}|g_1|^2 dx d\tau \right. \\ & \left. + s^2\lambda^2 \int_Q e^{2s\alpha}\varphi^2|g_2|^2 dx d\tau + s^3\lambda^4 \int_0^T \int_\omega e^{2s\alpha}\varphi^3|\xi|^2 dx d\tau \right). \end{aligned} \tag{4.25}$$

Finally, to get an estimate of $\nabla\xi$ we multiply by $s\lambda^2 e^{2s\alpha}\varphi\xi$ the equation (4.7) satisfied by ξ and we integrate by parts to obtain

$$\begin{aligned} & s\lambda^2 \int_Q e^{2s\alpha}\varphi|\nabla\xi|^2 dx d\tau + s\lambda^2 \int_Q e^{2s\alpha}\varphi\gamma|\xi_n|^2 dx d\tau = \\ & + s\lambda^2 \int_Q e^{2s\alpha}\varphi g_1 \cdot \xi dx d\tau + s\lambda^2 \int_Q \nabla(e^{2s\alpha}\varphi) \times \xi \cdot g_1 dx d\tau \\ & + s\lambda^2 \int_Q e^{2s\alpha}\varphi g_2 \cdot \text{curl} \xi dx d\tau - \frac{1}{2}s\lambda^2 \int_Q (e^{2s\alpha}\varphi)_t |\xi|^2 dx d\tau \\ & - s\lambda^2 \int_Q (\xi^t \nabla(e^{2s\alpha}\varphi)) : \nabla\xi dx d\tau. \end{aligned}$$

Then since for $s \geq C(T^k + T^{k-1})$ and $C > 0$ large enough we have $s\lambda^2 |(e^{2s\alpha}\varphi)_t| \leq Cs^3\lambda^2 e^{2s\alpha}\varphi^3$ and $s\lambda^2 |\nabla(e^{2s\alpha}\varphi)| \leq Cs^2\lambda^3 e^{2s\alpha}\varphi^2$, we deduce (4.18) from (4.25). \square

5. Proofs of Theorem 3.3 and of Theorem 3.4

5.1. Null controllability of the linear system: proof of Theorem 3.3

Assume $(g_1, g_2) \in \mathcal{G}$ and $(w_0, \theta_0) \in \mathbf{V}_0^{1/2}(\Omega) \times \widehat{\mathbf{V}}_n^{1/2}(\Omega)$ and recall first that according to Definition 6.12 of the appendix, the pair $(w, \theta) \in \mathbf{L}^2(Q) \times \mathbf{L}^2(Q)$ is a transposition solution of (1.19) if the following equality holds:

$$\begin{aligned} & \int_Q (w \cdot \mathcal{L}_1^*(y, \pi, \rho) + \theta \cdot \mathcal{L}_2^*(y, \kappa, \rho, \mu)) dx d\tau = \int_0^T \langle (g_1, g_2) | (y, \rho) \rangle_{\mathcal{V}, \mathcal{V}'} d\tau \\ & + \int_0^T \int_{\mathcal{O}} (u_1 \cdot y + u_2 \cdot P_{\mathcal{O}}\rho) dx d\tau + \int_{\Omega} (w_0 \cdot y(0) + \theta_0 \cdot \rho(0)) dx \end{aligned}$$

for all $(y, \pi, \rho, \kappa, \mu) \in \mathcal{R}_0$ with \mathcal{R}_0 and \mathcal{L}_i^* , $i = 1, 2$, defined in (6.21), (6.22) in appendix and $\mathcal{V} \stackrel{\text{def}}{=} \mathbf{V}_0^1(\Omega) \times \widehat{\mathbf{V}}_n^1(\Omega)$.

Thus, we define the following symmetric bilinear form on \mathcal{R}_0 :

$$\begin{aligned} b((y, \pi, \rho, \kappa, \mu), (y', \pi', \rho', \kappa', \mu')) &= \int_Q e^{s\frac{3\beta}{2}} (\mathcal{L}_1^*(y, \pi, \rho) \cdot \mathcal{L}_1^*(y', \pi', \rho')) dx d\tau \\ &+ \int_Q e^{s\frac{3\beta}{2}} \mathcal{L}_2^*(y, \kappa, \rho, \mu) \cdot \mathcal{L}_2^*(y', \kappa', \rho', \mu') dx d\tau \\ &+ \int_0^T \int_{\mathcal{O}} e^{s\frac{3\beta}{2}} (y \cdot y' + P_{\mathcal{O}}\rho \cdot P_{\mathcal{O}}\rho') dx d\tau, \end{aligned}$$

and the following linear form on \mathcal{R}_0 :

$$\begin{aligned} l((y', \pi', \rho', \kappa', \mu')) &= \int_0^T \langle (g_1, g_2) | (y', \rho') \rangle_{\mathcal{V}', \mathcal{V}} d\tau \\ &+ \int_{\Omega} w_0 \cdot y'(0) dx + \int_{\Omega} \theta_0 \cdot \rho'(0) dx. \end{aligned}$$

Inequality (3.12) guarantees that the semi norm induced by $b(\cdot, \cdot)$ is in fact a norm. Then we denote by \mathcal{R} the completion of \mathcal{R}_0 for this norm and, since $e^{-s\hat{\beta}}(g_1, g_2) \in L^2(0, T; \mathbf{V}_0^{-1}(\Omega) \times [\mathbf{H}_n^1(\Omega)]')$ and $(w_0, \theta_0) \in \mathbf{V}_n^0(\Omega) \times \widehat{\mathbf{V}}_n^0(\Omega)$, one also has that l is continuous on \mathcal{R} . Then by invoking Lax-Milgram Lemma we obtain a unique $(y, \pi, \rho, \kappa, \mu) \in \mathcal{R}$ satisfying for all $(y', \pi', \rho', \kappa', \mu') \in \mathcal{R}$:

$$b((y, \pi, \rho, \kappa, \mu), (y', \pi', \rho', \kappa', \mu')) = l((y', \pi', \rho', \kappa', \mu')). \quad (5.1)$$

Moreover, we have $\|(y, \pi, \rho, \kappa, \mu)\|_{\mathcal{R}} \leq C\|l\|_{\mathcal{R}'}$ or equivalently:

$$\begin{aligned} & \int_Q e^{s\frac{3\hat{\beta}}{2}} (|\mathcal{L}_1^*(y, \pi, \rho)|^2 + |\mathcal{L}_2^*(y, \kappa, \rho, \mu)|^2) dx d\tau \\ & + \int_0^T \int_{\mathcal{O}} e^{s\frac{3\hat{\beta}}{2}} (|y|^2 + |P_{\mathcal{O}}\rho|^2) dx d\tau \leq \\ & C \left(\|(g_1, g_2)\|_{\mathcal{G}}^2 + \|(w_0, \theta_0)\|_{\mathbf{V}_n^0(\Omega) \times \widehat{\mathbf{V}}_n^0(\Omega)}^2 \right). \end{aligned} \quad (5.2)$$

Thus, if we set $(w, \theta) = (e^{s\frac{3\hat{\beta}}{2}} \mathcal{L}_1^*(y, \pi, \rho), e^{s\frac{3\hat{\beta}}{2}} \mathcal{L}_2^*(y, \kappa, \rho, \mu))$ and $(u_1, u_2) = (-e^{s\frac{3\hat{\beta}}{2}} y, -e^{s\frac{3\hat{\beta}}{2}} P_{\mathcal{O}}\rho)$ then (5.1) implies that (w, θ) is a transposition solution of (1.19) and (5.2) yields

$$\begin{aligned} & \int_Q e^{-s\frac{3\hat{\beta}}{2}} (|w|^2 + |\theta|^2) dx d\tau + \int_0^T \int_{\mathcal{O}} e^{-s\frac{3\hat{\beta}}{2}} (|u_1|^2 + |u_2|^2) dx d\tau \leq \\ & C \left(\|(g_1, g_2)\|_{\mathcal{G}}^2 + \|(w_0, \theta_0)\|_{\mathbf{V}_n^0(\Omega) \times \widehat{\mathbf{V}}_n^0(\Omega)}^2 \right). \end{aligned} \quad (5.3)$$

Finally, since $(e^{-s\frac{\hat{\beta}}{2}} w, e^{-s\frac{\hat{\beta}}{2}} \theta)$ satisfies (1.19) with nonhomogeneous terms

$$\begin{aligned} & e^{-s\frac{\hat{\beta}}{2}} g_1 - s \frac{\widehat{\beta}_t}{2} e^{-s\frac{\hat{\beta}}{2}} w + e^{-s\frac{\hat{\beta}}{2}} \mathbf{1}_{\mathcal{O}} u_1 \in L^2(0, T; \mathbf{H}^{-1/2}(\Omega)), \\ & e^{-s\frac{\hat{\beta}}{2}} g_2 - s \frac{\widehat{\beta}_t}{2} e^{-s\frac{\hat{\beta}}{2}} \theta + e^{-s\frac{\hat{\beta}}{2}} \mathbf{1}_{\mathcal{O}} P_{\mathcal{O}} u_2 \in L^2(0, T; [\mathbf{H}_n^{1/2}(\Omega)]'), \end{aligned}$$

in transposition sense, and since by Proposition 6.13 in Appendix we have that transposition and weak solutions of (1.19) coincide, then by combining (5.3) with Theorem 6.9 in Appendix we get the estimate in Theorem 3.3.

5.2. Null controllability of the nonlinear system: proof of Theorem 3.4

The first part of the theorem will follow from the Banach fixed point theorem applied to the mapping $\Psi : (\widehat{w}, \widehat{\theta}) \mapsto (w, \theta)$ where (w, θ) is the solution given by Theorem 3.3 of the control problem (1.19) for $g_1 = g_1(\widehat{w}, \widehat{\theta})$ and $g_2 = g_2(\widehat{w}, \widehat{\theta})$ where

$$g_1(w, \theta) \stackrel{\text{def}}{=} (\theta \cdot \nabla) \theta - (w \cdot \nabla) w \quad \text{and} \quad g_2(w, \theta) \stackrel{\text{def}}{=} \text{curl}(w \times \theta).$$

For that, we first need some estimates for the nonlinear terms $g_1(w, \theta)$ and $g_2(w, \theta)$. From $\text{curl}(w \times \theta) = (\theta \cdot \nabla) w - (w \cdot \nabla) \theta$ and from classical estimates

of Navier-Stokes type nonlinearity one has:

$$\begin{aligned} \|g_1(w, \theta)\|_{\mathbf{H}^{-1/2}(\Omega)} &\leq C(\|\theta\|_{\mathbf{H}^{1/2}(\Omega)}\|w\|_{\mathbf{H}^{3/2}(\Omega)} + \|w\|_{\mathbf{H}^{1/2}(\Omega)}\|w\|_{\mathbf{H}^{3/2}(\Omega)}) \\ \|g_2(w, \theta)\|_{[\mathbf{H}_n^{1/2}(\Omega)]'} &\leq C(\|\theta\|_{\mathbf{H}^{1/2}(\Omega)}\|w\|_{\mathbf{H}^{3/2}(\Omega)} + \|w\|_{\mathbf{H}^{1/2}(\Omega)}\|\theta\|_{\mathbf{H}^{3/2}(\Omega)}). \end{aligned}$$

Then we deduce that

$$\begin{aligned} &\|e^{-s\hat{\beta}}(g_1(w, \theta), g_2(w, \theta))\|_{L^2(0, T; \mathbf{H}^{-1/2}(\Omega) \times [\mathbf{H}_n^{1/2}(\Omega)]')} \leq \\ &C(\|e^{-s\hat{\beta}/2}(w, \theta)\|_{L^\infty(0, T; \mathbf{V}_0^{1/2}(\Omega) \times \widehat{\mathbf{V}}_n^{1/2}(\Omega))} \|e^{-s\hat{\beta}/2}(w, \theta)\|_{L^2(0, T; \mathbf{V}_0^{3/2}(\Omega) \times \widehat{\mathbf{V}}_n^{3/2}(\Omega))}, \end{aligned}$$

and the continuous embedding $\mathcal{W} \hookrightarrow L^\infty(0, T; \mathbf{V}_0^{1/2}(\Omega) \times \widehat{\mathbf{V}}_n^{1/2}(\Omega))$ yields

$$\|(g_1(w, \theta), g_2(w, \theta))\|_{\mathcal{G}} \leq C\|(w, \theta)\|_{\mathcal{W}}^2. \quad (5.4)$$

Moreover, from analogous calculations we deduce that

$$\begin{aligned} &\|(g_1(w_1, \theta_1), g_2(w_1, \theta_1)) - (g_1(w_2, \theta_2), g_2(w_2, \theta_2))\|_{\mathcal{G}} \leq \\ &C(\|(w_1, \theta_1)\|_{\mathcal{W}} + \|(w_2, \theta_2)\|_{\mathcal{W}})\|(w_1 - w_2, \theta_1 - \theta_2)\|_{\mathcal{W}}. \end{aligned} \quad (5.5)$$

Thus, by combining (5.4), (5.5) and Theorem 3.3 we deduce that

$$\begin{aligned} &\|\Psi(w, \theta)\|_{\mathcal{W}} \leq c(\|(w, \theta)\|_{\mathcal{W}}^2 + \|(w_0, \theta_0)\|_{\mathbf{V}_0^{1/2}(\Omega) \times \widehat{\mathbf{V}}_n^{1/2}(\Omega)}), \\ &\|\Psi(w_1, \theta_1) - \Psi(w_2, \theta_2)\|_{\mathcal{W}} \leq c(\|(w_1, \theta_1)\|_{\mathcal{W}} + \|(w_2, \theta_2)\|_{\mathcal{W}}) \\ &\quad \times \|(w_1 - w_2, \theta_1 - \theta_2)\|_{\mathcal{W}}, \end{aligned}$$

for some constant $c > 0$ independent on $w, \theta, w_i, \theta_i, i = 1, 2$ and of (w_0, θ_0) . Then if we set $B_0 \stackrel{\text{def}}{=} \{(w, \theta) \in \mathcal{W} \mid \|(w, \theta)\|_{\mathcal{W}} \leq \delta\| (w_0, \theta_0)\|_{\mathbf{V}_0^{1/2}(\Omega) \times \widehat{\mathbf{V}}_n^{1/2}(\Omega)}\}$ one verifies that if $\|(w_0, \theta_0)\|_{\mathbf{V}_0^{1/2}(\Omega) \times \widehat{\mathbf{V}}_n^{1/2}(\Omega)} \leq \varepsilon$ and if $\delta > 0$ is large enough and $\varepsilon > 0$ is small enough such that $c(\delta^2\varepsilon + 1) < \delta$ and $2c\delta\varepsilon < 1$, the mapping Ψ is a contraction of B_0 into itself and (1.17)-(1.18) admits a unique solution in B_0 . Then the first part of Theorem 3.4 is proved.

The second part follows from a classical argument that we recall for the sake of completeness. If $(\tilde{w}, \tilde{\theta})$ is another solution of (1.17) in $C(0, T; \mathbf{H}^{1/2}(\Omega) \times \mathbf{H}^{1/2}(\Omega)) \cap L^2(0, T; \mathbf{H}^{3/2}(\Omega) \times \mathbf{H}^{3/2}(\Omega))$ then $(z, \vartheta) = (w - \tilde{w}, \theta - \tilde{\theta})$ satisfies

$$\left\{ \begin{array}{ll} z_t - \Delta z + (z \cdot \nabla)(\bar{v} + w) + ((\bar{v} + \tilde{w}) \cdot \nabla)z \\ \quad - ((\bar{B} + \theta) \cdot \nabla)\vartheta - (\vartheta \cdot \nabla)(\bar{B} + \tilde{\theta}) + \nabla\pi = 0 & \text{in } Q, \\ \vartheta_t + \text{curl}(\text{curl } \vartheta) - ((\bar{B} + \tilde{\theta}) \cdot \nabla)z + (z \cdot \nabla)(\bar{B} + \theta) \\ \quad - (\vartheta \cdot \nabla)(\bar{v} + w) + ((\bar{v} + \tilde{w}) \cdot \nabla)\vartheta = 0 & \text{in } Q, \\ \text{div } z = \text{div } \vartheta = 0 & \text{in } Q, \\ z = 0, \quad \vartheta \cdot n = 0 & \text{on } \Sigma, \\ (\text{curl } \vartheta - \bar{v} \times \vartheta) \times n = 0 & \text{on } \Sigma, \\ z(0) = 0, \quad \vartheta(0) = 0 & \text{in } \Omega. \end{array} \right. \quad (5.6)$$

To obtain (5.6) we have used the formula $\text{curl}(a \times b) = (b \cdot \nabla)a - (a \cdot \nabla)b$ for divergence free vector fields a, b . Then by multiplying the first equality

by z and the second equality by ϑ , by integrating over Ω and by making an integration by parts we obtain:

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} (\|z\|_{\mathbf{L}^2(\Omega)}^2 + \|\vartheta\|_{\mathbf{L}^2(\Omega)}^2) + \|\nabla z\|_{\mathbf{L}^2(\Omega)}^2 + \|\operatorname{curl} \vartheta\|_{\mathbf{L}^2(\Omega)}^2 &= - \int_{\Omega} (\bar{v} \cdot \nabla) \vartheta \cdot \vartheta dx \\ &\quad \int_{\Omega} (\vartheta \cdot \nabla) (\bar{v} + \tilde{w}) \cdot \vartheta dx - \int_{\Omega} (z \cdot \nabla) (\bar{v} + w) \cdot z dx - \int_{\Omega} (D^a(\bar{B} + \tilde{\theta})z \cdot \vartheta \cdot dx. \end{aligned}$$

Thus, from Sobolev embeddings we have the estimates:

$$\begin{aligned} \left| \int_{\Omega} (\bar{v} \cdot \nabla) \vartheta \cdot \vartheta dx \right| &\leq C \|\bar{v}\|_{\mathbf{H}^2(\Omega)} \|\vartheta\|_{\mathbf{H}^1(\Omega)} \|\vartheta\|_{\mathbf{L}^2(\Omega)}, \\ \left| \int_{\Omega} (\vartheta \cdot \nabla) (\bar{v} + \tilde{w}) \cdot \vartheta dx \right| &\leq C \|\bar{v} + \tilde{w}\|_{\mathbf{H}^{3/2}(\Omega)} \|\vartheta\|_{\mathbf{H}^{1/2}(\Omega)}^2, \\ \left| \int_{\Omega} (z \cdot \nabla) (\bar{v} + w) \cdot z dx \right| &\leq C \|\bar{v} + w\|_{\mathbf{H}^{3/2}(\Omega)} \|z\|_{\mathbf{H}^{1/2}(\Omega)}^2, \\ \left| \int_{\Omega} D^a(\bar{B} + \tilde{\theta})z \cdot \vartheta dx \right| &\leq C \|\bar{B} + \tilde{\theta}\|_{\mathbf{H}^{3/2}(\Omega)} \|z\|_{\mathbf{H}^{1/2}(\Omega)} \|\vartheta\|_{\mathbf{H}^{1/2}(\Omega)}, \end{aligned}$$

and the use of interpolation inequality $\|\cdot\|_{\mathbf{H}^{1/2}(\Omega)} \leq C \|\cdot\|_{\mathbf{H}^1(\Omega)}^{1/2} \|\cdot\|_{\mathbf{L}^2(\Omega)}^{1/2}$, Cauchy Schwartz inequality, (2.2) and Poincaré inequality, yields

$$\begin{aligned} \frac{d}{dt} (\|z\|_{\mathbf{L}^2(\Omega)}^2 + \|\vartheta\|_{\mathbf{L}^2(\Omega)}^2) + \|\nabla z\|_{\mathbf{L}^2(\Omega)}^2 + \|\operatorname{curl} \vartheta\|_{\mathbf{L}^2(\Omega)}^2 &\leq \\ C (\|\bar{v}\|_{\mathbf{H}^2(\Omega)}^2 + \|\bar{B}\|_{\mathbf{H}^2(\Omega)}^2 + \|w\|_{\mathbf{H}^{3/2}(\Omega)}^2 + \|\tilde{w}\|_{\mathbf{H}^{3/2}(\Omega)}^2) & \\ \times (\|z\|_{\mathbf{L}^2(\Omega)}^2 + \|\vartheta\|_{\mathbf{L}^2(\Omega)}^2). & \end{aligned}$$

Finally, by Gronwall's Lemma we get that z and ϑ are identically zero.

6. Appendix

6.1. Regularity results for Stokes problem with non standard conditions

The present section is dedicated to regularity results for the following Stokes system with a nonhomogeneous non standard boundary condition:

$$\left\{ \begin{array}{ll} \theta_t + \operatorname{curl}(\operatorname{curl} \theta) + \nabla \chi + \sum_{i=1}^N \beta_i \mathbf{g}_i = g & \text{in } Q, \\ \operatorname{div} \theta = 0 & \text{in } Q, \\ \forall i = 1, \dots, N \quad \int_{\Omega} \theta \cdot \mathbf{g}_i dx = 0 & \text{in } (0, T), \\ \theta \cdot n = 0, \quad \operatorname{curl} \theta \times n = b \times n & \text{on } \Sigma, \\ \theta(0) = \theta_0 & \text{in } \Omega. \end{array} \right. \quad (6.1)$$

Here, for $g \in \mathbf{L}^2(Q)$ and $\theta_0 \in \widehat{\mathbf{V}}_n^1(\Omega)$ we aim at providing regularity results for a boundary datum $b \in \mathbf{H}^{\frac{r}{2}, \frac{r}{4}}(\Sigma)$ for $r \in [0, 1]$.

But before to deal with system (6.1), we need preliminary lemmas.

Lemma 6.1. For $f \in \mathbf{L}^2(\Omega)$ satisfying $\int_{\Omega} f \cdot g_i dx = 0$, $i = 1, \dots, N$ and $h \in H_0^1(\Omega)$ satisfying $\int_{\Omega} h dx = 0$ the following Stokes system with non standard boundary conditions

$$\begin{cases} \operatorname{curl}(\operatorname{curl} \theta) + \nabla \chi = f & \text{in } \Omega, \\ \operatorname{div} \theta = h & \text{in } \Omega, \\ \theta \cdot n = 0 & \text{on } \partial\Omega, \\ \operatorname{curl} \theta \times n = 0 & \text{on } \partial\Omega, \end{cases} \quad (6.2)$$

admits a unique solution $(\theta, \chi) \in \mathbf{H}^2(\Omega) \times H^1(\Omega)$ satisfying $\int_{\Omega} \theta \cdot g_i dx = 0$ for all $i = 1, \dots, N$, and $\int_{\Omega} \chi dx = 0$. Moreover, the following estimate holds

$$\|\theta\|_{\mathbf{H}^2(\Omega)} + \|\chi\|_{H^1(\Omega)} \leq C(\|f\|_{\mathbf{L}^2(\Omega)} + \|h\|_{H^1(\Omega)})$$

for a constant $C > 0$ independent on θ, χ, f, h .

Proof. Let us first note that we can reduce the problem to the case $h = 0$ with a lifting procedure. Indeed, since $h \in H_0^1(\Omega)$ and $\int_{\Omega} h dx = 0$ there exists $\theta_h \in \mathbf{H}_0^2(\Omega)$ such that $\operatorname{div} \theta_h = h$ in Ω and $\|\theta_h\|_{\mathbf{H}^2(\Omega)} \leq C\|h\|_{H^1(\Omega)}$ for some $C > 0$ independent on h and θ_h , see [2, Cor. 3.1]. Then θ is a solution of (6.2) if and only if $\theta - \theta_h$ satisfies (6.2) with $h = 0$ and with a nonhomogeneous body force term $f - \operatorname{curl}(\operatorname{curl} \theta_h)$ which depends continuously on $(f, h) \in \mathbf{L}^2(\Omega) \times H^1(\Omega)$.

It remains to consider the system (6.2) with an homogeneous divergence condition, namely:

$$\begin{cases} \operatorname{curl}(\operatorname{curl} \theta) + \nabla \chi = f & \text{in } \Omega, \\ \operatorname{div} \theta = 0 & \text{in } \Omega, \\ \theta \cdot n = 0 & \text{on } \partial\Omega, \\ \operatorname{curl} \theta \times n = 0 & \text{on } \partial\Omega. \end{cases} \quad (6.3)$$

By multiplying the first equation of (6.3) by a test function $\rho \in \widehat{\mathbf{V}}_n^1(\Omega)$ we obtain the following weak formulation of (6.3):

$$\forall \rho \in \widehat{\mathbf{V}}_n^1(\Omega) \quad \int_{\Omega} \operatorname{curl} \theta \cdot \operatorname{curl} \rho dx = \int_{\Omega} f \cdot \rho dx. \quad (6.4)$$

By (2.2) we have that the bilinear form defined at the left of the above equality is coercive on $\widehat{\mathbf{V}}_n^1(\Omega)$ and the Lax-Milgram Lemma yields the existence and the uniqueness of $\theta \in \widehat{\mathbf{V}}_n^1(\Omega)$ solution to the variational problem (6.4). Moreover, since $\operatorname{curl} g_i = 0$ in Ω and $\int_{\Omega} f \cdot g_i dx = 0$ for $i = 1, \dots, N$ the formulation (6.4) is also true for test functions ρ in $\mathbf{V}_n^1(\Omega)$. We denote by $P : \mathbf{L}^2(\Omega) \rightarrow \mathbf{V}_n^0(\Omega)$ the Helmholtz orthogonal projection operator related to Ω . From the underlying related Neumann problem (see [25, Chap.III, Lem 1.2]) we can verify that P is also continuous from $\mathbf{H}^1(\Omega)$ onto $\mathbf{V}_n^1(\Omega)$. Then for all $\rho \in \mathbf{H}^1(\Omega)$ we have $P\rho \in \mathbf{V}_n^1(\Omega)$ and

$$\forall \rho \in \mathbf{H}^1(\Omega) \quad \int_{\Omega} \operatorname{curl} \theta \cdot \operatorname{curl} P\rho dx = \int_{\Omega} f \cdot P\rho dx, \quad (6.5)$$

and since $\operatorname{curl} P\rho = \operatorname{curl} \rho$ in Ω we deduce that

$$\forall \rho \in \mathbf{H}^1(\Omega) \quad \int_{\Omega} \operatorname{curl} \theta \cdot \operatorname{curl} \rho dx = \int_{\Omega} (f - \nabla \chi) \cdot \rho dx, \quad (6.6)$$

where $\chi \in H^1(\Omega)$ is given by $\nabla \chi = (I - P)f$. It follows that $\operatorname{curl}(\operatorname{curl} \theta) = f - \nabla \chi \in \mathbf{L}^2(\Omega)$ which allows to define $\operatorname{curl} \theta \times n$ in $\mathbf{H}^{-\frac{1}{2}}(\partial\Omega)$ from the following formula for all $\rho \in \mathbf{H}^1(\Omega)$:

$$\langle (\operatorname{curl} \theta \times n), \rho \rangle_{\mathbf{H}^{-\frac{1}{2}}(\partial\Omega), \mathbf{H}^{\frac{1}{2}}(\partial\Omega)} = \int_{\Omega} \operatorname{curl} \theta \cdot \operatorname{curl} \rho dx - \int_{\Omega} \operatorname{curl}(\operatorname{curl} \theta) \cdot \rho dx.$$

Then from (6.6) we deduce that $\operatorname{curl} \theta \times n = 0$ on $\partial\Omega$. Finally, since $z \stackrel{\text{def}}{=} \operatorname{curl} \theta$ satisfies $\operatorname{curl} z \in \mathbf{L}^2(\Omega)$, $\operatorname{div} z = 0$ in Ω and $z \times n = 0$ on $\partial\Omega$ we first get $\operatorname{curl} \theta \in \mathbf{H}^1(\Omega)$ from the first statement of [1, Cor. 2.15]. Thus, from $\operatorname{curl} \theta \in \mathbf{H}^1(\Omega)$, $\operatorname{div} \theta = 0$ in Ω and $\theta \cdot n = 0$ on $\partial\Omega$ which is of class $C^{2,1}$, we finally get $\theta \in \mathbf{H}^2(\Omega)$ from the second statement of [1, Cor. 2.15]. \square

The above result permits to deduce the following lifting Lemma.

Lemma 6.2. *For $b \in \mathbf{H}^{\frac{1}{2}, \frac{1}{4}}(\Sigma)$ there exists $\theta_b \in \mathbf{H}^{2,1}(Q)$ satisfying*

$$\theta_b(0) = 0 \text{ in } \Omega, \quad \operatorname{div} \theta_b = 0 \text{ in } Q, \quad (6.7)$$

$$\operatorname{curl} \theta_b \times n = b \times n \quad \text{and} \quad \theta_b \cdot n = 0 \text{ on } \Sigma. \quad (6.8)$$

Moreover, the following estimate holds

$$\|\theta_b\|_{\mathbf{H}^{2,1}(Q)} \leq c \|b\|_{\mathbf{H}^{\frac{1}{2}, \frac{1}{4}}(\Sigma)} \quad (6.9)$$

for some constant $c > 0$ independent on b .

Proof. We first invoke [27, Thm 7.2] to obtain the existence of $\vartheta_b \in \mathbf{H}^{2,1}(Q)$ depending continuously on $b \in \mathbf{H}^{\frac{1}{2}, \frac{1}{4}}(\Sigma)$ and such that $\vartheta_b(0) = 0$ in Ω , $\vartheta_b = 0$ on Σ and $\frac{\partial \vartheta_b}{\partial n} = b \times n$ on Σ . Note that since $\vartheta_b = 0$ on Σ implies $\operatorname{curl} \vartheta_b \times n = \frac{\partial \vartheta_b}{\partial n}$ on Σ we deduce that $\operatorname{curl} \vartheta_b \times n = b \times n$ on Σ . Thus, we set $\theta_b = \vartheta_b - L\vartheta_b$ where the linear mapping L is defined by $L\vartheta = \theta \in \widehat{\mathbf{V}}_n^2(\Omega)$ where θ is the solution given by Lemma 6.1 of the Stokes problem:

$$\begin{cases} \operatorname{curl}(\operatorname{curl} \theta) + \nabla \chi = 0 & \text{in } \Omega, \\ \operatorname{div} \theta = \operatorname{div} \vartheta & \text{in } \Omega, \\ \theta \cdot n = 0 & \text{on } \partial\Omega, \\ \operatorname{curl} \theta \times n = 0 & \text{on } \partial\Omega. \end{cases} \quad (6.10)$$

It is clear that such a θ_b will satisfies (6.7) and (6.8). According to Lemma 6.1, L is continuous from $\{\vartheta \in \mathbf{H}^2(\Omega) \cap \mathbf{H}_0^1(\Omega) \mid \operatorname{div} \vartheta = 0 \text{ on } \partial\Omega\}$ into $\mathbf{H}^2(\Omega)$. Then since we have $\vartheta_b(t) = 0$ on $\partial\Omega$ and $\operatorname{div} \vartheta_b(t) = \frac{\partial \vartheta_b(t)}{\partial n} \cdot n = 0$ on $\partial\Omega$, we deduce that $L\vartheta_b(t)$ is well defined and that $\|(I - L)\vartheta_b\|_{L^2(0,T;\mathbf{H}^2(\Omega))} \leq C \|\vartheta_b\|_{L^2(0,T;\mathbf{H}^2(\Omega))}$. Moreover, with a transposition argument we can also

prove that L is continuous for the norm of $\mathbf{L}^2(\Omega)$: if we multiply the first equation of (6.10) by ρ solution of

$$\begin{cases} \operatorname{curl}(\operatorname{curl} \rho) + \nabla \kappa = \theta & \text{in } \Omega, \\ \operatorname{div} \rho = 0 & \text{in } \Omega, \\ \rho \cdot n = 0 & \text{on } \partial\Omega, \\ \operatorname{curl} \rho \times n = 0 & \text{on } \partial\Omega, \end{cases} \quad (6.11)$$

an integration by parts gives $\int_{\Omega} |\theta|^2 dx = \int_{\Omega} \vartheta \cdot \nabla \kappa dx$. Thus, by using the bound $\|\nabla \kappa\|_{\mathbf{L}^2(\Omega)} \leq C\|\theta\|_{\mathbf{L}^2(\Omega)}$ given by Lemma 6.1 we get $\|L\vartheta\|_{\mathbf{L}^2(\Omega)} \leq C\|\vartheta\|_{\mathbf{L}^2(\Omega)}$ and then $\|(I - L)\vartheta_b\|_{H^1(0,T;\mathbf{L}^2(\Omega))} \leq C\|\vartheta_b\|_{H^1(0,T;\mathbf{L}^2(\Omega))}$. Finally, we have $\|\theta_b\|_{\mathbf{H}^{2,1}(Q)} \leq C\|\vartheta_b\|_{\mathbf{H}^{2,1}(Q)}$ and the conclusion follows from the continuous dependence of θ_b with respect to b . \square

Finally, we are now in position to turn back to system 6.1.

Definition 6.3. For $\theta_0 \in \widehat{\mathbf{V}}_n^1(\Omega)$, $g \in \mathbf{L}^2(Q)$ and $b \in \mathbf{H}^{\frac{r}{2}, \frac{r}{4}}(\Sigma)$ with $r \in [0, 1]$, the function $\theta \in W(0, T; \widehat{\mathbf{V}}_n^1(\Omega), [\widehat{\mathbf{V}}_n^1(\Omega)]')$ is said to be a weak solution of (6.1) if and only if $\theta(0) = \theta_0$ and if for all $\rho \in \widehat{\mathbf{V}}_n^1(\Omega)$ and $t \in (0, T)$ we have:

$$\frac{d}{dt} \int_{\Omega} \theta(t) \cdot \rho dx + \int_{\Omega} \operatorname{curl} \theta(t) \cdot \operatorname{curl} \rho dx = \int_{\Omega} g(t) \cdot \rho dx + \int_{\partial\Omega} (b(t) \times n) \cdot \rho d\Gamma.$$

Remark 6.4. As usual, the above equality is obtained from (6.1) by (formally) multiplying the first equality by ρ and integrating by parts. Conversely, if θ is a regular weak solution, for instance $\theta \in W(0, T; \widehat{\mathbf{V}}_n^2(\Omega), \widehat{\mathbf{V}}_n^0(\Omega))$, then an integration by parts and the use of (2.3) permits to recover the formulation (6.1) in a classical sense for some $\chi \in L^2(0, T; H^1(\Omega))$ and $\beta_i \in L^2(0, T)$.

Theorem 6.5. Assume $r \in [0, 1]$ and $\epsilon \in (0, \frac{1}{2})$. For all $\theta_0 \in \widehat{\mathbf{V}}_n^1(\Omega)$, $g \in \mathbf{L}^2(Q)$ and $b \in \mathbf{H}^{\frac{r}{2}, \frac{r}{4}}(\Sigma)$ the system (6.1) admits a unique weak solution θ . Moreover, $\theta \in \mathbf{H}^{\frac{3}{2} - \epsilon + (\frac{1}{2} + \epsilon)r, \frac{3}{4} - \frac{\epsilon}{2} + (\frac{1}{4} + \frac{\epsilon}{2})r}(Q)$ and satisfies:

$$\|\theta\|_{\mathbf{H}^{\frac{3}{2} - \epsilon + (\frac{1}{2} + \epsilon)r, \frac{3}{4} - \frac{\epsilon}{2} + (\frac{1}{4} + \frac{\epsilon}{2})r}(Q)} \leq c(\|b\|_{\mathbf{H}^{\frac{r}{2}, \frac{r}{4}}(\Sigma)} + \|g\|_{\mathbf{L}^2(Q)} + \|\theta_0\|_{\mathbf{V}_n^1(\Omega)}), \quad (6.12)$$

for some constant $c > 0$ independent on θ , b , θ_0 , g .

Proof. The weak formulation in Definition 6.3 writes

$$\theta' + \widehat{A}\theta = \widehat{P}g + Tb,$$

where \widehat{A} is the linear operator defined from the bilinear form $(\theta, \rho) \in \widehat{\mathbf{V}}_n^1(\Omega) \times \widehat{\mathbf{V}}_n^1(\Omega) \mapsto \int_{\Omega} \operatorname{curl} \theta \cdot \operatorname{curl} \rho dx$, where \widehat{P} is the orthogonal projector from $\mathbf{L}^2(\Omega)$ onto $\widehat{\mathbf{V}}_n^0(\Omega)$ and where $Tb \in [\widehat{\mathbf{V}}_n^1(\Omega)]'$ is defined by $\langle Tb | \rho \rangle = \int_{\partial\Omega} b \times n \cdot \rho d\Gamma$. Classical arguments guarantee that \widehat{A} generates an analytic semigroup on $\mathbf{V}_n^0(\Omega)$ (see [9, Thm. 2.12, Part II, Chap 1]) and from Lemma 6.1 we deduce that $D(\widehat{A}) = \{\theta \in \mathbf{V}_n^2(\Omega) \mid \operatorname{curl} \theta \times n = 0 \text{ on } \partial\Omega\}$. Moreover, since we easily verify that $Tb \in [\widehat{\mathbf{V}}_n^{\frac{1}{2} + \epsilon}(\Omega)]'$, maximal regularity results with an interpolation

argument give $\theta \in W(0, T; \widehat{\mathbf{V}}_n^{\frac{3}{2}-\epsilon}(\Omega), [\widehat{\mathbf{V}}_n^{\frac{1}{2}+\epsilon}(\Omega)]')$. Then (6.12) for $r = 0$ follows because this last space is continuously embedded in $\mathbf{H}^{\frac{3}{2}-\epsilon, \frac{3}{4}-\frac{\epsilon}{2}}(Q)$.

It remains to prove (6.12) for $r = 1$, the case $r \in (0, 1)$ will then follow by interpolation. According to Lemma 6.2 there exists $\theta_b \in \mathbf{H}^{2,1}(Q)$ satisfying (6.7), (6.8) and depending continuously on $b \in \mathbf{H}^{\frac{1}{2}, \frac{1}{4}}(\Sigma)$. Then $\vartheta = \theta - \theta_b$ satisfies

$$\vartheta' + \widehat{A}\vartheta = \widehat{P}\tilde{g}, \quad \vartheta(0) = \theta_0$$

for $\tilde{g} = g - (\theta_b)_t - \text{curl}(\text{curl } \theta_b) \in \mathbf{L}^2(Q)$, and (6.12) for $r = 1$ follows from classical maximal regularity results for analytic semigroup and from the continuity of $b \in \mathbf{H}^{\frac{1}{2}, \frac{1}{4}}(\Sigma) \mapsto \theta_b \in \mathbf{H}^{2,1}(Q)$. \square

6.2. Regularity results for linear MHD equations

The present subsection deals with the well-posedness of the following non autonomous linear magnetohydrodynamic equations:

$$\left\{ \begin{array}{ll} w_t - \Delta w + (w \cdot \nabla)\bar{v} + (\bar{v} \cdot \nabla)w - (\bar{B} \cdot \nabla)\theta \\ \quad - (\theta \cdot \nabla)\bar{B} + \nabla q = g_1 & \text{in } Q, \\ \theta_t + \text{curl}(\text{curl } \theta) - \text{curl}(w \times \bar{B} + \bar{v} \times \theta) \\ \quad + \nabla \chi + \sum_{i=1}^N \beta_i g_i = g_2 & \text{in } Q, \\ \text{div } w = \text{div } \theta = 0 & \text{in } Q, \\ \forall i = 1, \dots, N \quad \int_{\Omega} \theta \cdot g_i dx = 0 & \text{in } (0, T), \\ w = 0, \quad \theta \cdot n = 0 & \text{on } \Sigma, \\ (\text{curl } \theta - \bar{v} \times \theta) \times n = 0 & \text{on } \Sigma, \\ w(0) = w_0, \quad \theta(0) = \theta_0 & \text{in } \Omega. \end{array} \right. \quad (6.13)$$

For the following we assume $(g_1, g_2) \in L^2(0, T; \mathbf{H}^{-1}(\Omega)) \times [\mathbf{H}_n^1(\Omega)]'$ and $(w_0, \theta_0) \in \mathbf{V}_n^0(\Omega) \times \widehat{\mathbf{V}}_n^0(\Omega)$. In (6.13) the pressure functions q, χ and the real values $\beta_i, i = 1, \dots, N$ play the role of the Lagrange multipliers related to constraints $\text{div } w = \text{div } \theta = 0$ in Q and $\int_{\Omega} \theta \cdot g_i dx = 0, i = 1, \dots, N$ in $(0, T)$.

Remark 6.6. If g_2 is of the form $\text{curl}(w \times \theta) + \mathbf{1}_{\mathcal{O}} P_{\mathcal{O}} u_2$ then we have $\nabla \chi = 0$ in Q and $\beta_i = 0, i = 1, \dots, N$, and the induction equation in (1.19) is recovered. This can be obtained by successively multiplying the equation by $\nabla \chi$ and g_i , integrating by parts and using (1.11), $g_i \in \mathbf{V}_n^0(\Omega)$ and $\text{curl } g_i = 0$ in Ω .

We first introduce the notion of weak solution of (6.13). For that we set

$$\mathcal{H} \stackrel{\text{def}}{=} \mathbf{V}_n^0(\Omega) \times \widehat{\mathbf{V}}_n^0(\Omega) \quad \text{and} \quad \mathcal{V} \stackrel{\text{def}}{=} \mathbf{V}_0^1(\Omega) \times \widehat{\mathbf{V}}_n^1(\Omega)$$

and we define the family of linear operators $A(t) : \mathcal{V} \rightarrow \mathcal{V}'$, $t \in [0, T]$ by

$$\begin{aligned} \langle A(t)(w, \theta) | (y, \rho) \rangle_{\mathcal{V}, \mathcal{V}'} &\stackrel{\text{def}}{=} \int_{\Omega} (\nabla w : \nabla y + \text{curl } \theta \cdot \text{curl } \rho) dx \\ &+ \int_{\Omega} ((w \cdot \nabla) \bar{v}(t) + (\bar{v}(t) \cdot \nabla) w - (\bar{B}(t) \cdot \nabla) \theta - (\theta \cdot \nabla) \bar{B}(t)) \cdot y dx \\ &- \int_{\Omega} (w \times \bar{B}(t) + \bar{v}(t) \times \theta) \cdot \text{curl } \rho dx. \end{aligned}$$

Definition 6.7. Assume $(g_1, g_2) \in L^2(0, T; \mathbf{H}^{-1}(\Omega) \times [\mathbf{H}_n^1(\Omega)]')$ and $(w_0, \theta_0) \in \mathcal{H}$. We say that $(w, \theta) \in W(0, T; \mathcal{V}, \mathcal{V}')$ is a weak solution of (6.13) if it satisfies for all $(y, \rho) \in \mathcal{V}$ and $t \in (0, T)$:

$$\begin{aligned} \frac{d}{dt} \langle (w(t), \theta(t)) | (y, \rho) \rangle_{\mathcal{H}} \\ + \langle A(t)(w(t), \theta(t)) | (y, \rho) \rangle_{\mathcal{V}, \mathcal{V}'} = \langle (g_1(t), g_2(t)) | (y, \rho) \rangle_{\mathcal{V}, \mathcal{V}'} \quad (6.14) \\ (w(0), \theta(0)) = (w_0, \theta_0). \end{aligned}$$

Remark 6.8. Such a notion of weak solution is coherent because if (g_1, g_2) and (w, θ) are regular, for instance $(g_1, g_2) \in \mathbf{L}^2(Q) \times \mathbf{L}^2(Q)$ and $(w, \theta) \in W(0, T; \mathbf{V}_n^2(\Omega) \times \widehat{\mathbf{V}}_n^2(\Omega), \mathbf{V}_n^0(\Omega) \times \widehat{\mathbf{V}}_n^0(\Omega))$, then (w, θ) satisfies (6.13) in a classical sense. This can be deduced from an integration by parts and by recalling (2.1) and (2.3) to obtain the existence of $q \in L^2(0, T; H^1(\Omega))$, $\chi \in L^2(0, T; H^1(\Omega))$ and $\beta_i \in L^2(0, T)$, $i = 1, \dots, N$, such that (6.13) is satisfied.

We have the following existence theorem for weak solutions.

Theorem 6.9. Assume $r \in [0, 1]$, $(w_0, \theta_0) \in \mathbf{V}_0^r(\Omega) \times \widehat{\mathbf{V}}_n^r(\Omega)$ and $(g_1, g_2) \in L^2(0, T; \mathbf{H}^{r-1}(\Omega) \times [\mathbf{H}_n^{1-r}(\Omega)]')$. Then the system (6.13) admits a unique weak solution (w, θ) . Moreover, $(w, \theta) \in W(0, T; \mathbf{V}_0^{r+1}(\Omega) \times \widehat{\mathbf{V}}_n^{r+1}(\Omega), \mathbf{V}^{r-1}(\Omega) \times [\widehat{\mathbf{V}}_n^{1-r}(\Omega)]')$ and satisfies:

$$\begin{aligned} \|(w, \theta)\|_{W(0, T; \mathbf{V}_0^{r+1}(\Omega) \times \widehat{\mathbf{V}}_n^{r+1}(\Omega), \mathbf{V}^{r-1}(\Omega) \times [\widehat{\mathbf{V}}_n^{1-r}(\Omega)]')} &\leq \\ C \|(g_1, g_2)\|_{L^2(0, T; \mathbf{H}^{r-1}(\Omega) \times [\mathbf{H}_n^{1-r}(\Omega)]')} + \|(w_0, \theta_0)\|_{\mathbf{V}_0^r(\Omega) \times \widehat{\mathbf{V}}_n^r(\Omega)}, \quad (6.15) \end{aligned}$$

for a constant $C > 0$ independent on w, θ and on w_0, θ_0, g_1, g_2 .

Proof. From (1.8) and standard Sobolev embeddings we get that for all $(w, \theta) \in \mathcal{V}$ and $(y, \rho) \in \mathcal{V}$ the mapping $t \mapsto \langle (A(t))(w, \theta) | (y, \rho) \rangle_{\mathcal{V}, \mathcal{V}'}$ is measurable on $(0, T)$ and that the following continuity and coercivity inequalities hold for all $t \in (0, T)$:

$$\begin{aligned} \langle A(t)(w, \theta) | (y, \rho) \rangle_{\mathcal{V}, \mathcal{V}'} &\leq c_1 \|(w, \theta)\|_{\mathcal{V}} \|(y, \rho)\|_{\mathcal{V}}, \\ \langle A(t)(w, \theta) | (w, \theta) \rangle_{\mathcal{V}, \mathcal{V}'} + c_2 \|(w, \theta)\|_{\mathcal{H}}^2 &\geq c_3 \|(w, \theta)\|_{\mathcal{V}}^2, \end{aligned}$$

for some $c_1 > 0$, $c_2 > 0$ and $c_3 > 0$ independent on t and (w, θ, y, ρ) . Then according to [9, Part II, Chap. 2, Thm 1.2], for $(g_1, g_2) \in L^2(0, T; \mathbf{H}^{-1}(\Omega) \times [\mathbf{H}_n^1(\Omega)]')$ and $(w_0, \theta_0) \in \mathcal{H}$ the equations (6.14) admits a unique solution satisfying (6.15) for $r = 0$.

It remains to prove (6.15) for $r = 1$. Then the conclusion will follow by interpolation. For the following we suppose that $(g_1, g_2) \in \mathbf{L}^2(Q) \times \mathbf{L}^2(Q)$ and $(w_0, \theta_0) \in \mathbf{V}_n^1(\Omega) \times \widehat{\mathbf{V}}_n^1(\Omega)$. From (6.14) we have that w is a weak solution of the following Stokes system:

$$\begin{cases} w_t - \Delta w + \nabla q = \tilde{g}_1 & \text{in } Q, \\ \operatorname{div} w = 0 & \text{in } Q, \\ w = 0 & \text{on } \Sigma, \\ w(0) = w_0 & \text{in } \Omega, \end{cases} \quad (6.16)$$

with $\tilde{g}_1 \stackrel{\text{def}}{=} g_1 - (w \cdot \nabla) \bar{v} - (\bar{v} \cdot \nabla) w + (\bar{B} \cdot \nabla) \theta + (\theta \cdot \nabla) \bar{B} \in \mathbf{L}^2(Q)$, and we have that θ is a weak solution of the following Stokes system with non standard boundary conditions:

$$\begin{cases} \theta_t + \operatorname{curl}(\operatorname{curl} \theta) + \nabla \chi + \sum_{i=1}^N \beta_i g_i = \tilde{g}_2 & \text{in } Q, \\ \operatorname{div} \theta = 0 & \text{in } Q, \\ \forall i = 1, \dots, N \quad \int_{\Omega} \theta \cdot g_i dx = 0 & \text{in } (0, T), \\ \theta \cdot n = 0, \quad \operatorname{curl} \theta \times n = b \times n & \text{on } \Sigma, \\ \theta(0) = \theta_0 & \text{in } \Omega, \end{cases} \quad (6.17)$$

where $\tilde{g}_2 = g_2 + \operatorname{curl}(w \times \bar{B} + \bar{v} \times \theta) \in \mathbf{L}^2(Q)$ and $b = \bar{v} \times \theta$. Since $\tilde{g}_1 \in \mathbf{L}^2(Q)$ classical regularity results for the Stokes problem with homogeneous boundary conditions yield $w \in W(0, T; \mathbf{V}_n^2(\Omega) \times \mathbf{V}_n^0(\Omega))$. It remains to prove that $\theta \in W(0, T; \widehat{\mathbf{V}}_n^2(\Omega), \widehat{\mathbf{V}}_n^0(\Omega))$ and for that we need to combine regularity results for system (6.17) of Theorem 6.5 with a bootstrap argument.

First, note that (1.8) implies in particular that

$$\bar{v}|_{\Sigma} \in L^\infty(0, T; \mathbf{H}^{\frac{1}{2}}(\partial\Omega)) \cap H^{\frac{1}{2}}(0, T; \mathbf{H}^{\frac{1}{2}}(\partial\Omega)). \quad (6.18)$$

Then since θ given by (6.14) belongs to $W(0, T; \mathbf{V}_n^1(\Omega), [\widehat{\mathbf{V}}_n^1(\Omega)]')$ we have $\theta \in \mathbf{H}^{1, \frac{1}{2}}(Q)$ and $\theta|_{\Sigma} \in \mathbf{H}^{\frac{1}{2}, \frac{1}{4}}(\Sigma)$, and with (6.18) classical Sobolev embeddings guarantee that $\bar{v} \times \theta|_{\Sigma} \in \mathbf{L}^2(\Sigma)$. Thus, from (6.12) we get that $\theta \in \mathbf{H}^{\frac{3}{2}-\epsilon, \frac{3}{4}-\frac{\epsilon}{2}}(Q)$ and then $\theta|_{\Sigma} \in \mathbf{H}^{1-\epsilon, \frac{1}{2}-\frac{\epsilon}{2}}(\Sigma)$ for $\epsilon \in (0, \frac{1}{4})$. Then with (6.18) we get $\bar{v} \times \theta|_{\Sigma} \in \mathbf{H}^{\frac{1}{2}-2\epsilon, \frac{1}{4}-\epsilon}(\Sigma)$ and (6.12) yields $\theta \in \mathbf{H}^{2-2\epsilon, 1-\epsilon}(Q)$. It follows that $\theta|_{\Sigma} \in \mathbf{H}^{\frac{3}{2}-2\epsilon, \frac{3}{4}-\epsilon}(\Sigma)$ and using again (6.18) we deduce that $\bar{v} \times \theta|_{\Sigma} \in \mathbf{H}^{\frac{1}{2}, \frac{1}{4}}(\Sigma)$. Finally, from (6.12) we get $\theta \in \mathbf{H}^{2,1}(Q)$ and then (6.15) holds for $r = 1$. This ends the proof. \square

Let us now consider the well-posedness of the adjoint system:

$$\left\{ \begin{array}{ll} -y_t - \Delta y - (D^s y)\bar{v} + (D^a \rho)\bar{B} + \nabla \pi = f_1 & \text{in } Q, \\ -\rho_t - \Delta \rho + (D^s y)\bar{B} - (D^a \rho)\bar{v} + \nabla \kappa + \sum_{i=1}^N \mu_i g_i = f_2 & \text{in } Q, \\ \operatorname{div} y = \operatorname{div} \rho = 0 & \text{in } Q, \\ \forall i = 1, \dots, N \int_{\Omega} \rho \cdot g_i dx = 0 & \text{in } (0, T), \\ y = 0, \quad \rho \cdot n = 0, \quad (\operatorname{curl} \rho) \times n = 0 & \text{on } \Sigma. \\ y(T) = y_T \quad \rho(T) = \rho_T & \text{in } \Omega. \end{array} \right. \quad (6.19)$$

Similarly as for system (6.13) we define weak solutions of (6.19) as follows.

Definition 6.10. Assume that $(f_1, f_2) \in L^2(0, T; \mathbf{H}^{-1}(\Omega)) \times [\mathbf{H}_n^1(\Omega)]'$ and that $(y_T, \rho_T) \in \mathcal{H}$. We say that $(y, \rho) \in W(0, T; \mathcal{V}, \mathcal{V}')$ is a weak solution of (6.13) if it satisfies for all $(w, \theta) \in \mathcal{V}$ and $t \in (0, T)$:

$$\begin{aligned} & -\frac{d}{dt} \langle (w, \theta) | (y(t), \rho(t)) \rangle_{\mathcal{H}} \\ & + \langle A(t)(w, \theta) | (y(t), \rho(t)) \rangle_{\mathcal{V}, \mathcal{V}'} = \langle (f_1(t), f_2(t)) | (w, \theta) \rangle_{\mathcal{V}, \mathcal{V}'} \\ & (y(T), \rho(T)) = (y_T, \rho_T). \end{aligned}$$

We have the following regularity theorem for weak solutions of (6.13).

Theorem 6.11. Assume $(y_T, \rho_T) \in \mathbf{V}_0^1(\Omega) \times \widehat{\mathbf{V}}_n^1(\Omega)$ and $(f_1, f_2) \in \mathbf{L}^2(Q) \times \mathbf{L}^2(Q)$. Then system (6.19) admits a unique weak solution (y, ρ) . Moreover, (y, ρ) belongs to $W(0, T; \mathbf{V}_0^2(\Omega) \times \widehat{\mathbf{V}}_n^2(\Omega), \mathbf{V}_n^0(\Omega) \times \widehat{\mathbf{V}}_n^0(\Omega))$ and satisfies:

$$\begin{aligned} & \|(y, \rho)\|_{W(0, T; \mathbf{V}_0^2(\Omega) \times \widehat{\mathbf{V}}_n^2(\Omega), \mathbf{V}_n^0(\Omega) \times \widehat{\mathbf{V}}_n^0(\Omega))} \leq \\ & C(\|(f_1, f_2)\|_{\mathbf{L}^2(Q) \times \mathbf{L}^2(Q)} + \|(y_T, \rho_T)\|_{\mathbf{V}_0^1(\Omega) \times \widehat{\mathbf{V}}_n^1(\Omega)}) \end{aligned} \quad (6.20)$$

for a constant $C > 0$ independent on y, ρ and on y_T, ρ_T, f_1, f_2 .

Since the proof of Theorem 6.11 is similar to that of Theorem 6.9 we omit it.

Finally, we introduce the notion of transposition solutions of (6.13) which is equivalent to the notion of weak solution and which is required for the duality argument in Section 5. For that we introduce the space

$$\begin{aligned} \mathcal{R}_0 \stackrel{\text{def}}{=} & \{ (y, \pi, \rho, \kappa, \mu) \in \mathbf{C}^2(\overline{Q}) \times C^1(\overline{Q}) \times \mathbf{C}^2(\overline{Q}) \times C^1(\overline{Q}) \times C([0, T])^N \mid \\ & y = 0, \quad \rho \cdot n = 0 \text{ and } \operatorname{curl} \rho \times n = 0 \text{ on } \Sigma, \\ & \operatorname{div} y = \operatorname{div} \rho = 0 \text{ in } Q, \\ & \forall t \in [0, T] \int_{\Omega} \rho(t) \cdot g_i dx = 0, \quad i = 1, \dots, N, \\ & \forall t \in [0, T] \int_{\Omega} \pi(t) dx = \int_{\Omega} \kappa(t) dx = 0 \}, \end{aligned} \quad (6.21)$$

and for $(y, \pi, \rho, \kappa, \mu) \in \mathcal{R}_0$ we set:

$$\begin{aligned} \mathcal{L}_1^*(y, \pi, \rho) &\stackrel{\text{def}}{=} -y_t - \Delta y - (D^s y)\bar{v} + (D^a \rho)\bar{B} + \nabla \pi, \\ \mathcal{L}_2^*(y, \kappa, \rho, \mu) &\stackrel{\text{def}}{=} -\rho_t - \Delta \rho + (D^s y)\bar{B} - (D^a \rho)\bar{v} + \nabla \kappa + \sum_{i=1}^N \mu_i g_i. \end{aligned} \quad (6.22)$$

Definition 6.12. Assume that $(w_0, \theta_0) \in \mathbf{V}_n^0(\Omega) \times \widehat{\mathbf{V}}_n^0(\Omega)$ and that $(g_1, g_2) \in L^2(0, T; \mathbf{H}^{-1}(\Omega)) \times [\mathbf{H}_n^1(\Omega)]'$. We say that $(w, \theta) \in \mathbf{L}^2(Q) \times \mathbf{L}^2(Q)$ is a transposition solution of (1.19) if for all $(y, \pi, \rho, \kappa, \mu) \in \mathcal{R}_0$ such that $(y(T), \rho(T)) = (0, 0)$ the following equality holds:

$$\begin{aligned} &\int_Q (w \cdot \mathcal{L}_1^*(y, \pi, \rho) + \theta \cdot \mathcal{L}_2^*(y, \kappa, \rho, \mu)) dx d\tau \\ &= \int_0^T \langle (g_1, g_2) | (y, \rho) \rangle_{\mathcal{V}', \mathcal{V}} d\tau + \int_\Omega (w_0 \cdot y(0) + \theta_0 \cdot \rho(0)) dx. \end{aligned} \quad (6.23)$$

Proposition 6.13. For $(w_0, \theta_0) \in \mathbf{V}_n^0(\Omega) \times \widehat{\mathbf{V}}_n^0(\Omega)$ and $(g_1, g_2) \in L^2(0, T; \mathbf{H}^{-1}(\Omega)) \times [\mathbf{H}_n^1(\Omega)]'$, (w, θ) is a weak solution of (1.19) if and only if it is a transposition solution of (1.19).

Proof. We first remark that the existence and the uniqueness of a transposition solution (w, θ) is a direct consequence of Theorem 6.11. Indeed, since a density argument guarantees that the transposition solution (w, θ) must satisfy for all $(f_1, f_2) \in \mathbf{L}^2(Q) \times \mathbf{L}^2(Q)$

$$\int_Q (w \cdot f_1 + \theta \cdot f_2) dx d\tau = \int_0^T \langle (g_1, g_2) | (y, \rho) \rangle_{\mathcal{V}', \mathcal{V}} d\tau + \int_\Omega (w_0 \cdot y(0) + \theta_0 \cdot \rho(0)) dx,$$

where $(y, \rho) \in W(0, T; \mathbf{V}_0^2(\Omega) \times \widehat{\mathbf{V}}_n^2(\Omega), \mathbf{V}_n^0(\Omega) \times \widehat{\mathbf{V}}_n^0(\Omega))$ is the corresponding solution of (6.19) with $(y_T, \rho_T) = (0, 0)$, the conclusion follows from the Riesz representation theorem. Then it follows that weak solution and transposition solution coincide. Indeed, to get that every weak solution is a transposition solution it suffices to choose a smooth (y, ρ) in formulation (6.14) and to integrate by parts to recover formulation (6.23). Thus, the fact that every transposition solution is a weak solution follows from the uniqueness of the transposition solution. \square

References

- [1] C. Amrouche, C. Bernardi, M. Dauge, and V. Girault. Vector potentials in three-dimensional non-smooth domains. *Math. Methods Appl. Sci.*, 21(9):823–864, 1998.
- [2] C. Amrouche and V. Girault. Decomposition of vector spaces and application to the Stokes problem in arbitrary dimension. *Czechoslovak Math. J.*, 44(119)(1):109–140, 1994.
- [3] M. Badra. Global Carleman inequalities for Stokes and penalized Stokes equations. *Math. Control Relat. Fields*, 1(2):149–175, 2011.

- [4] M. Badra, S. Ervedoza, and S. Guerrero. Local exact controllability for the density-dependent incompressible Navier-Stokes equations.
- [5] M. Badra and T. Takahashi. On Fattorini criterion for approximate controllability and stabilizability of parabolic equations. preprint.
- [6] V. Barbu, T. Havârneanu, C. Popa, and S. S. Sritharan. Exact controllability for the magnetohydrodynamic equations. *Comm. Pure Appl. Math.*, 56(6):732–783, 2003.
- [7] V. Barbu, T. Havârneanu, C. Popa, and S. S. Sritharan. Local exact controllability for the magnetohydrodynamic equations revisited. *Adv. Differential Equations*, 10(5):481–504, 2005.
- [8] V. Barbu, S. S. Rodrigues, and A. Shirikyan. Internal exponential stabilization to a nonstationary solution for 3D Navier-Stokes equations. *SIAM J. Control Optim.*, 49(4):1454–1478, 2011.
- [9] A. Bensoussan, G. Da Prato, M. C. Delfour, and S. K. Mitter. *Representation and control of infinite dimensional systems*. Systems & Control: Foundations & Applications Birkhäuser Boston Inc., Boston, MA, second edition, 2007.
- [10] J.-M. Coron and A. V. Fursikov. Global exact controllability of the 2D Navier-Stokes equations on a manifold without boundary. *Russian J. Math. Phys.*, 4(4):429–448, 1996.
- [11] J.-M. Coron and S. Guerrero. Null controllability of the N -dimensional Stokes system with $N - 1$ scalar controls. *J. Differential Equations*, 246(7):2908–2921, 2009.
- [12] R. Dautray and J.-L. Lions. *Analyse mathématique et calcul numérique pour les sciences et les techniques. Vol. 5*. INSTN: Collection Enseignement. [INSTN: Teaching Collection]. Masson, Paris, 1988. Spectre des opérateurs. [The operator spectrum], With the collaboration of Michel Artola, Michel Cessenat, Jean Michel Combes and Bruno Scheurer, Reprinted from the 1984 edition.
- [13] P. A. Davidson. *An introduction to magnetohydrodynamics*. Cambridge Texts in Applied Mathematics. Cambridge University Press, Cambridge, 2001.
- [14] E. Fernández-Cara, M. González-Burgos, S. Guerrero, and J.-P. Puel. Null controllability of the heat equation with boundary Fourier conditions: the linear case. *ESAIM Control Optim. Calc. Var.*, 12(3):442–465 (electronic), 2006.
- [15] E. Fernández-Cara and S. Guerrero. Global Carleman inequalities for parabolic systems and applications to controllability. *SIAM J. Control Optim.*, 45(4):1399–1446 (electronic), 2006.
- [16] E. Fernández-Cara and S. Guerrero. Local exact controllability of micropolar fluids. *J. Math. Fluid Mech.*, 9(3):419–453, 2007.
- [17] E. Fernández-Cara, S. Guerrero, O. Yu. Imanuvilov, and J.-P. Puel. Local exact controllability of the Navier-Stokes system. *J. Math. Pures Appl. (9)*, 83(12):1501–1542, 2004.
- [18] E. Fernández-Cara, S. Guerrero, O. Yu. Imanuvilov, and J.-P. Puel. Some controllability results for the N -dimensional Navier-Stokes and Boussinesq systems with $N - 1$ scalar controls. *SIAM J. Control Optim.*, 45(1):146–173 (electronic), 2006.
- [19] C. Foias̃ and R. Temam. Remarques sur les équations de Navier-Stokes stationnaires et les phénomènes successifs de bifurcation. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 5(1):28–63, 1978.

- [20] A. V. Fursikov. Exact boundary zero controllability of three-dimensional Navier-Stokes equations. *J. Dynam. Control Systems*, 1(3):325–350, 1995.
- [21] A. V. Fursikov and O. Yu. Èmanuilov. Exact controllability of the Navier-Stokes and Boussinesq equations. *Uspekhi Mat. Nauk*, 54(3(327)):93–146, 1999.
- [22] A. V. Fursikov and O. Yu. Imanuvilov. On exact boundary zero-controllability of two-dimensional Navier-Stokes equations. *Acta Appl. Math.*, 37(1-2):67–76, 1994. Mathematical problems for Navier-Stokes equations (Centro, 1993).
- [23] A. V. Fursikov and O. Yu. Imanuvilov. On controllability of certain systems simulating a fluid flow. In *Flow control (Minneapolis, MN, 1992)*, volume 68 of *IMA Vol. Math. Appl.*, pages 149–184. Springer, New York, 1995.
- [24] A. V. Fursikov and O. Yu. Imanuvilov. *Controllability of evolution equations*, volume 34 of *Lecture Notes Series*. Seoul National University Research Institute of Mathematics Global Analysis Research Center, Seoul, 1996.
- [25] G. P. Galdi. *An introduction to the mathematical theory of the Navier-Stokes equations. Vol. I. Linearized steady problems*, volume 38 of *Springer Tracts in Natural Philosophy*. Springer-Verlag, New York, 1994.
- [26] M. González-Burgos, S. Guerrero, and J.-P. Puel. Local exact controllability to the trajectories of the Boussinesq system via a fictitious control on the divergence equation. *Commun. Pure Appl. Anal.*, 8(1):311–333, 2009.
- [27] P. Grisvard. Caractérisation de quelques espaces d’interpolation. *Arch. Rational Mech. Anal.*, 25:40–63, 1967.
- [28] S. Guerrero. Local exact controllability to the trajectories of the Boussinesq system. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 23(1):29–61, 2006.
- [29] S. Guerrero. Controllability of systems of Stokes equations with one control force: existence of insensitizing controls. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 24(6):1029–1054, 2007.
- [30] S. Guerrero and F. Guillén-González. On the controllability of the hydrostatic Stokes equations. *J. Math. Fluid Mech.*, 10(3):402–422, 2008.
- [31] T. Havârneanu, C. Popa, and S. S. Sritharan. Exact internal controllability for the magnetohydrodynamic equations in multi-connected domains. *Adv. Differential Equations*, 11(8):893–929, 2006.
- [32] T. Havârneanu, C. Popa, and S. S. Sritharan. Exact internal controllability for the two-dimensional magnetohydrodynamic equations. *SIAM J. Control Optim.*, 46(5):1802–1830 (electronic), 2007.
- [33] O. Y. Imanuvilov and M. Yamamoto. Carleman inequalities for parabolic equations in Sobolev spaces of negative order and exact controllability for semilinear parabolic equations. *Publ. Res. Inst. Math. Sci.*, 39(2):227–274, 2003.
- [34] O. Yu. Imanuvilov. Remarks on exact controllability for the Navier-Stokes equations. *ESAIM Control Optim. Calc. Var.*, 6:39–72 (electronic), 2001.
- [35] O. Yu. Imanuvilov, J.-P. Puel, and M. Yamamoto. Carleman estimates for parabolic equations with nonhomogeneous boundary conditions. *Chin. Ann. Math. Ser. B*, 30(4):333–378, 2009.
- [36] O. Yu. Imanuvilov, J.-P. Puel, and M. Yamamoto. Carleman estimates for second order nonhomogeneous parabolic equations. Preprint.
- [37] O. Yu. Imanuvilov and T. Takahashi. Exact controllability of a fluid-rigid body system. *J. Math. Pures Appl. (9)*, 87(4):408–437, 2007.

- [38] T. Kim and D. Cao. Local exact controllability of the Navier-Stokes equations with the condition on the pressure on parts of the boundary. *SIAM J. Control Optim.*, 48(6):3805–3837, 2010.
- [39] C.-G. Lefter. On a unique continuation property related to the boundary stabilization of magnetohydrodynamic equations. *An. Ştiinţ. Univ. Al. I. Cuza Iaşi. Mat. (N.S.)*, 56(1):1–15, 2010.
- [40] J.-L. Lions and E. Magenes. *Problèmes aux limites non homogènes et applications. Vol. 1.* Travaux et Recherches Mathématiques, No. 17. Dunod, Paris, 1968.
- [41] J.-P. Raymond and M. Vanninathan. Null controllability in a fluid-solid structure model. *J. Differential Equations*, 248(7):1826–1865, 2010.
- [42] M. Tucsnak and G. Weiss. *Observation and control for operator semigroups.* Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks]. Birkhäuser Verlag, Basel, 2009.

Mehdi Badra
Laboratoire LMA, UMR CNRS 5142
Université de Pau et des Pays de l'Adour
64013 Pau Cedex
France
e-mail: mehdi.badra@univ-pau.fr